

509,009

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



24 SEP 2004



(43) International Publication Date  
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number  
**WO 03/080663 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/47**,  
C12N 15/12, C07K 16/18, A61K 38/17, C12N 15/11,  
5/10, G01N 33/50

(74) Common Representative: **NOVARTIS  
FORSCHUNGSSTIFTUNG, ZWEIGNIEDERLAS-  
SUNG FRIEDRICH MIESCHER INSTITUTE FOR  
BIOMEDICAL RESEARCH**; Maulbeerstrasse 66,  
CH-4058 Basel (CH).

(21) International Application Number: **PCT/EP03/03150**

(22) International Filing Date: 26 March 2003 (26.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0207224.7 27 March 2002 (27.03.2002) GB

(71) Applicant (for all designated States except US): **NOVAR-  
TIS FORSCHUNGSSTIFTUNG, ZWEIGNIEDER-  
LASSUNG FRIEDRICH MIESCHER INSTITUTE  
FOR BIOMEDICAL RESEARCH** [CH/CH]; Maulbeer-  
strasse 66, CH-4058 Basel (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MANN, Ruth** [CH/CH]; Gstaad, CH-4153  
Reinach (CH). **SCHERBERICH, Arnau** [FR/FR]; 1,  
rue des Ecrins, F-68400 Riedisheim (FR).

(54) Title: **TENASCIN-W COMPOSITIONS AND USES THEREOF**

(57) Abstract: Tenascin-W, an extracellular matrix molecule that is specifically expressed in metastatic tumours is provided. A system comprising a sample expressing tenascin-W is used as an *in vitro* method for screening possible anti-tumour agents or for agents that promote osteogenesis.

BEST AVAILABLE COPY



WO 03/080663 A2

TENASCIN-W COMPOSITIONS AND USES THEREOF

5 The present invention relates to polypeptides specifically expressed in tumours, to active agents having anti-tumour and/or anti-tumourigenic activity and to agents effective in improving conditions dependent on stem cell differentiation, in particular osteoblast formation, such as in osteogenesis, to pharmaceutical compositions of these agents and to the pharmaceutical uses  
10 of such agents and compositions. The invention also relates to *in vitro* methods of screening agents for anti-tumour and/or anti-tumourigenic activity as well as for agents effective in promoting stem cell differentiation.

The adherence of cells to each other and to the extracellular matrix (ECM) as  
15 well as the cellular signals transduced as a consequence of such binding are of fundamental importance to the development and maintenance of body form and function. The ECM has an important regulatory function in tissue homeostasis and, together with oncogenes and tumour suppressor genes is  
involved in tumourigenesis (reviewed in Boudreau, N. & Bissell, M. J.  
20 (1998) *Curr Opin Cell Biol* 10: 640-646 and Ruoslahti, E. (1999) *Adv Cancer Res* 76: 1-20).

In the more affluent countries of the world cancer is the cause of death of roughly one person in five with the five most common cancers being those of  
25 the lung, stomach, breast, colon/rectum and the uterine cervix. Tumors of this type often metastasize through lymphatic and vascular channels. Cancer is not fatal in every case and only about half the number of people who develop cancer die of it. The problem facing cancer patients and their physicians is that seeking to cure cancer is like trying to get rid of weeds.

30

One way to treat cancer effectively is to get an early diagnosis. Most cancers are not extensively vascularized (and therefore not invasive) during the early stages of development. The transition to a highly vascularized, invasive and ultimately metastatic cancer which spreads throughout the body commonly

takes ten years or longer. If the cancer is detected prior to invasion, surgical removal of the cancerous tissue is an effective cure. However, cancer is often detected only upon manifestation of clinical symptoms. Generally, such symptoms are present only when the disease is well established, often after metastasis has occurred, and the prognosis for the patient is poor, even after surgical resection of the cancerous tissue. Early detection of cancer therefore is important in that detection may significantly reduce morbidity. A reliable, non-invasive, and accurate technique for diagnosing cancer at an early stage would help save many lives.

10

Cancer cells can be removed surgically or destroyed with toxic compounds or with radiation but it is very hard to eliminate all of the cancerous cells. A general goal is therefore to find better ways of selectively killing cancer cells whilst leaving normal cells of the body unaffected. Part of that effort involves identifying new anti-cancer agents.

15

Apart from tumorigenesis, the ECM has an important regulatory function in tissue homeostasis and in the development and maintenance of body form and function, e.g. in the development or remodeling of skeleton or in bone morphogenesis. Bone marrow has stem cells with osteogenic potential and is made up of determined osteogenic precursor cells that are committed to osteogenesis and of inducible osteogenic precursor cells. Determined osteogenic precursor cells can differentiate into bone without an exogenous signal. Inducible osteogenic precursor cells require a molecular signal for initiating the differentiation program, e.g. induction by binding to extracellular matrix.

20

25

A number of molecules mediating cell adhesion have been identified and characterized at the molecular level both in vertebrates and invertebrates. Tenascins are a family of large multimeric extracellular matrix proteins, each having homologous subunits built from variable numbers of repeated domains. These include heptad repeats, epidermal growth factor (EGF)-like repeats, fibronectin type III domains and a C-terminal globular domain which is also found in fibrinogens. Tenascin-C was the first member of the family to

30

be discovered, in one instance as a myotendinous antigen (Chiquet, M. & Fambrough, DM. (1984) J Cell Biol 98(6):1937-1946) and in another, as a protein enriched in the stroma of gliomas (Bourdon, MA. et al (1983) Cancer Res 43(6):2796-2805, reflecting the major sites of tenascin-C expression, namely in tendons and ligaments and the extracellular matrix of tumor stroma. A further instance of the discovery of tenascin-C (also termed hexabrachion) reflects its interaction with fibronectin (Erickson, HP. et al. (1984) Nature 311(5983):267-9). Enforced interaction of tumour cells with fibronectin can block proliferation in cell culture and can decrease tumour growth in nude mice (Akamatsu H. et al (1996) Cancer Res 56: 4541-4546 and Giancotti, F. G & Ruoslahti, E. (1990) Cell 60: 849-859). Tenascin-C was shown to disrupt the interaction of cells with fibronectin and in this manner may enhance tumour cell proliferation. Chiquet-Ehrismann, R. et al (1988) Cell 53: 383-390 were the first to show that tenascin-C binds to fibronectin, blocks cell attachment to fibronectin and increases proliferation of rat breast adenocarcinoma cells (Chiquet-Ehrismann, R. et al (1988) Cell 53: 131-139).

Tenascin-C is present in a large number of developing tissues including the nervous system. Although abundant in mature ligaments and tendons, it is absent from skeletal and heart muscle, unless the muscle has been injured. Tenascin-C expression is elevated in essentially all carcinomas as well as in many other types of tumors (for review see Chiquet-Ehrismann, R. (1993) Semin Cancer Biol 4(5):301-10). Furthermore, tenascin-C is upregulated in wound healing (Latijnhouwers, MA. et al. (1996) J Pathol 178(1):30-5), during skeletogenesis (Koyama, E. et al (1996) J Orthop Res. 14(3):403-412 and Hall, BK. & Miyake, T. (1995) Int J Dev Biol. 39(6):881-893) as well as in many diseases involving infections and inflammation (Schenk, S. et al. (1995) Int J Cancer 61(4):443-9).

Each tenascin family member exhibits a specific gene expression pattern during embryogenesis and in the adult (for review see Chiquet-Ehrismann, R. (1995) Experientia 51(9-10):853-62) suggesting specific roles for each member. Tenascin-R is an extracellular matrix component of the nervous system found mainly in brain tissue (Pasheva, P. et al. (2001) Prog Brain Res.

132:103-14. Review), whereas tenascin-X is prominently expressed in muscle and skin connective tissue. In one patient, tenascin-X deficiency has been reported to result in an Ehler's Danlos phenotype (Burch, GH. et al. (1997) Nat Genet 17(1):104-8).

5

To date there is only one report on tenascin-W available in the literature. (Weber, P. et al. (1998) J Neurobiol 35(1):1-16). In this study, a cDNA encoding tenascin-W was isolated from a 20-28h postfertilization zebrafish cDNA library on the basis of the conserved epidermal growth factor-like domains found in all tenascin molecules. The expression pattern of tenascin-W transcripts was studied in the developing zebrafish by in situ hybridisation. It was found to be present in neural crest and sclerotome cells and the developing skeleton. Genebank sequence AJ001423 provides a zebrafish tenascin-W, and AL049689 provides a "novel human mRNA from chromosome 1, similar to Tenascin-R", whose function is not known.

10  
15

The present invention provides a composition comprising an isolated nucleic acid molecule having a nucleotide sequence selected from the group consisting of:

20

- (a) a nucleotide sequence as set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO: 2;
- (c) a nucleotide sequence with at least 85% identity to the sequence of (a) or (b);
- (d) a subsequence of more than 50 consecutive nucleotides of a sequence of (a), (b) or (c); and
- (e) a nucleotide sequence complementary to any of the nucleotide sequences or subsequence in (a), (b), (c) or (d).

25

30

In one aspect of the invention, the isolated nucleic acid molecule having a nucleotide sequence preferably exhibits at least 85% identity to the sequence of (a), more preferably encoding a variant of the amino acid sequence shown in SEQ ID NO: 2, such as a variant comprising an amino acid deletion, addition (e.g. fusion proteins) or substitution of the amino acid sequence

shown in SEQ ID NO:2. Preferably, the variant comprises a conservative substitution of at least one amino acid in said amino acid sequence in SEQ ID: NO: 2, more preferably the variant has stem cell differentiation inducing activity, in particular an activity that induces osteoblast development from stem cells. Most preferred is when the isolated nucleic acid molecule encodes a protein with the amino acid sequence shown in SEQ ID NO: 2.

The nucleic acid molecule can be an antisense molecule, in which case it might be desirable to have nucleotide residues that are resistant to nuclease degradation substituting some or all of the ribo- or deoxyribonucleotides.

Also provided are nucleic acid vectors comprising the nucleic acid molecules of the invention, as well as host cells comprising the vectors or nucleic acids, and transgenic, knockout or genetically modified animals (other than humans, in particular mice), comprising manipulated nucleic acids of the invention or absent or foreign sequence.

The invention also provides a composition comprising an isolated polypeptide having an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence as set forth in SEQ ID No. 2; and
- (b) an amino acid sequence with at least 85% identity to the sequence of (a); and
- (c) a subsequence of at least 30 consecutive amino acids of the sequence of (a) or (b), with the proviso that said subsequence does not fall within amino acid nos.1102 and 1152 of SEQ ID NO:2.

Preferably, the amino acid sequence in (b) comprises a conservative substitution of at least one amino acid of the amino acid sequence of SEQ ID: NO: 2. More preferably, the polypeptide or fragment has stem cell differentiation inducing activity, as described above. Useful fragments may exhibit an epitope recognized by polyclonal antibodies raised against the polypeptide having the amino acid sequence shown in SEQ ID NO: 2, for example. A particularly preferred polypeptide is that encoded by the amino acid sequence shown in SEQ ID NO: 2.

Also provided are antibodies that are specifically reactive against the polypeptides of the invention.

- 5 In another aspect of the invention, a composition comprising an isolated nucleic acid molecule having a nucleotide sequence selected from the group consisting of:
- (a) a nucleotide sequence as set forth in SEQ ID No. 1 or SEQ ID No. 3;
  - (b) a nucleotide sequence encoding the amino acid sequence shown in  
10 SEQ ID NO: 2 or SEQ ID NO: 4;
  - (c) a nucleotide sequence with at least 35% identity to any one of the sequences of (a) or (b), preferably (a);
  - (d) a subsequence of a least 15 consecutive nucleotides of the sequence of (a),(b) or (c); and
  - 15 (e) a nucleotide sequence complementary to (a),(b), (c), or (d), and a pharmaceutically acceptable excipient, diluent or carrier.

In one embodiment, the nucleic acid molecule preferably encodes a protein having stem cell differentiation inducing activity. In another embodiment, the  
20 nucleic acid molecule has a subsequence that is antisense to SEQ ID NO:1 or SEQ ID NO:3, wherein the nucleic acid molecule may comprise nucleotide residues that are resistant to nuclease degradation. In another embodiment, the isolated nucleic acid molecule encodes the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO:4. In yet another embodiment, the nucleic acid  
25 molecule has a subsequence selected from the group consisting of nucleotides 2380-3171 of SEQ ID No:1, nucleotides 2371-3162 of SEQ ID No:3, a complement of nucleotides 2380-3171 of SEQ ID No:1, and a complement of nucleotides 2371-3162 of SEQ ID No:3, or an RNA equivalent thereof.

30

Thus, also provided are nucleic acid compositions as described above for use as a pharmaceutical, as well as the use of such compositions for the manufacture of a medicament for the prophylaxis or treatment of cancer or bone pathologies.

Also provided are compositions comprising tenascin-W, preferably recombinant tenascin-W, and a pharmaceutically acceptable excipient, diluent or carrier. In preferred embodiments, the tenascin-W is a polypeptide having  
5 an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence as set forth in SEQ ID No. 2 or 4;
- (b) an amino acid sequence with at least 35% identity to the sequence of (a); and
- (c) a subsequence of at least 30 consecutive amino acids of the  
10 sequence of (a) or (b).

Preferably, the polypeptide has stem cell differentiation inducing activity as described above. More preferably, the polypeptide is encoded by the amino acid sequence shown in SEQ ID NO: 4.

15 Thus, also provided is the use of tenascin-W for the treatment of any disease or condition requiring increased bone mass, e.g. thrombosis, wound healing or atherosclerosis, as well as any condition ameliorated by the promotion of osteogenesis, e.g. bone healing,  
20 osteoporosis, as well as the use of tenascin-W as a stem cell marker.

Also provided are antibodies that specifically recognizes tenascin W for use as a pharmaceutical, as well as for the manufacture of a medicament, for the prophylaxis or treatment of cancer (e.g., glioblastoma, prostate, lung,  
25 colorectal, osteo- or breast carcinoma), including metastatic cancer, or for the prophylaxis or treatment of any disease or condition involving tenascin-W, e.g. excessive bone growth.

The present invention also provides methods for identifying agents for the  
30 prevention or the prophylactic treatment of tumourigenesis or the treatment or prophylactic treatment of tumours or cancer, or the treatment or prophylactic treatment of any disease or condition involving tenascin-W, e.g. a condition ameliorated by the promotion or inhibition of osteogenesis, comprising



contacting a test compound with a tenascin-W expressing cell sample and then measuring a change in one or more of:

- (a) cell proliferation, e.g. cell progression entering S-phase of the cell cycle;
  - 5 (b) DNA synthesis;
  - (c) cell adhesion;
  - (d) cell spreading;
  - (e) focal adhesion and actin stress fibre formation on fibronectin; and
  - (f) cell binding to extracellular matrix (ECM)
- 10 relative to when said test compound is absent.

Optionally, the method further comprises measuring a change in tenascin-W expression relative to when the test compound is absent. The tenascin-W may have any one or more of the features described above. A particularly  
15 preferred assays is carried out in the form of an enzyme linked immunosorbent assay (ELISA).

Also provided is a method for identifying regulators of tenascin W function, comprising:

- 20 (a) contacting a test compound with tenascin W and/or alpha8 beta1 integrin, and
  - (b) measuring the binding of the test compound to tenascin-W and/or alpha8 beta1 integrin, or
  - (c) measuring a disruption of binding of tenascin-W to alpha8 beta1  
25 integrin,
- relative to when the test compound is absent.

Optionally the method further comprises measuring the binding of a control compound to tenascin-W. In one embodiment, the tenascin-W is attached to a  
30 solid surface, for example using an antibody reactive against tenascin-W. The binding can be conveniently detected using an antibody labelled with a fluorescent label, a fluorescence quencher, a radioactive label, a scintillant or an enzyme. Alternatively, the binding is detected by measuring the adhesion of alpha8 beta1 to the immobilized tenascin-W (as described in example 8) or

vice versa. A decrease in binding of tenascin-W to alpha8 beta1 integrin is indicative of an inhibitor of the tenascin-W to alpha8 beta1 integrin interaction (and therefore an inhibitor of tenascin W function). An increase in binding of tenascin-W to alpha8 beta1 integrin in the presence of a test compound is indicative of a potential agent that activates alpha8 beta1 integrin, thereby acting as an agonist of tenascin-W function.

Thus also provided, are agents for the prevention or the prophylactic treatment of tumourigenesis or the diagnosis or the treatment or prophylactic treatment of tumours, or the treatment or prophylactic treatment of any disease or condition involving tenascin-W, e.g. a condition ameliorated by the promotion of osteogenesis, identified by a screening method of the invention.

Also provided are methods of diagnosing or prognosing cancer comprising:

- (a) analysing a sample obtained from an individual for the presence of tenascin-W
- (b) correlating the presence of tenascin-W with an unfavourable prognosis or diagnosis

Optionally, the method may further comprise correlating in an increase in (elevated level of) tenascin-W in the sample relative to healthy tissue with an unfavourable prognosis or diagnosis. Tenascin-W can be conveniently detected using an antibody specific for tenascin-W or alternatively tenascin-w can be detected at the transcript level using techniques well known in the art, such as a polymerase chain reaction (e.g., RT-PCR). The method may also include the additional use of controls.

The sample can be blood serum from an individual, for example. The method may also further comprise propagating cells in a sample in cell culture. In one embodiment, the method further comprises analysing the sample for the presence of alpha 8 integrin, the presence of alpha 8 integrin correlating with an unfavourable prognosis or diagnosis.

The present inventors have investigated extracellular matrix molecules, their  
5 expression during development, cell adhesion and proliferation of tumour cells  
and have characterized a novel member of the mammalian tenascin family.  
Prior to the present invention, no tenascin-W had been identified from a  
mammalian source and its function was not previously known. The present  
10 inventors have identified and characterized the complete cDNA sequence  
encoding the mouse and human tenascin-W. Antisera were prepared against  
a fragment of tenascin-W, which detect tenascin W in tumour stroma, in the  
periosteum and in liver tissue, and cross react with tenascin W from several  
mammalian species. In particular, the inventors have discovered that  
15 tenascin-W is specifically expressed in metastatic tumour cells as well as in  
the periosteum, the stem cell compartment for osteogenesis.

Thus, in one aspect, the present invention provides a composition comprising  
an isolated nucleic acid molecule having a nucleotide sequence selected from  
the group consisting of:

- 20 (a) a nucleotide sequence as set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence encoding the amino acid sequence  
shown in SEQ ID NO: 2;
- (c) a nucleotide sequence with at least 85% identity to the sequence  
of (a) or (b);
- 25 (d) a subsequence of more than 50, 75, 100, 200 or more  
consecutive nucleotides of a sequence of (a), (b) or (c); and
- (e) a nucleotide sequence complementary to any of the nucleotide  
sequences or subsequence in (a), (b), (c) or (d).

30 The compositions include various types of nucleic acid, including genomic  
DNA, cDNA and mRNA, for example. In one aspect of the invention, the  
isolated nucleic acid molecule having a nucleotide sequence preferably  
exhibits at least 85% identity, more preferably 90% identity, most preferably  
95, 98 or 100% identity to the sequence of (a) (SEQ ID NO:1). Also

encompassed are nucleic acids that encode polypeptides having the amino acid sequence shown in SEQ ID NO: 2, or variants thereof such as a variant comprising an amino acid deletion, addition (e.g. fusion proteins) or substitution relative to the amino acid sequence shown in SEQ ID NO:2. The various nucleic acids that can encode these polypeptides therefore may differ because of the degeneracy of the genetic code, in that most amino acids are encoded by more than one triplet codon. The identity of such codons is well known in this art, and this information can be used for the construction of the nucleic acids within the scope of the invention. Variants differ from wild-type protein in having one or more amino acid substitutions that either enhance, add, or diminish a biological activity of the wild-type protein. Once the amino acid change is selected, a nucleic acid encoding that variant is constructed according to methods well known in the art.

Preferably, the variant comprises a conservative substitution of at least one amino acid in said amino acid sequence in SEQ ID: NO: 2. The amino acids typically exhibit a biological function of the polypeptide as set forth in SEQ ID NO:2, that is, stem cell differentiation inducing activity; in particular, activity that induces osteoblast development from stem cells, or binding to an antibody that specifically recognizes Tenascin-W. To maintain biological activity, only conservative substitutions are therefore preferred as is well known in the art. Most preferred is when the isolated nucleic acid molecule encodes a protein with the amino acid sequence shown in SEQ ID NO: 2.

The nucleic acid molecule can be an antisense molecule, in which case it might be desirable to have nucleotide residues that are resistant to nuclease degradation substituting some or all of the ribo- or deoxyribonucleotides. Such nucleotide residues resistant to nucleases are well known in the art and can be easily synthesized by chemical means.

Also provided are nucleic acid vectors comprising the nucleic acid molecules of the invention, as well as host cells comprising the vectors or nucleic acids, and transgenic, knockout or genetically modified animals (other than humans,

in particular mice), comprising manipulated nucleic acids of the invention or absent the endogenous sequence.

The invention also provides a composition comprising an isolated polypeptide having an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence as set forth in SEQ ID NO: 2;
- (b) an amino acid sequence with at least 85% identity, preferably 90, 95, 98 or 100% identity to the sequence of (a); and
- (c) a subsequence of at least 30, 40, 50, 75, 100 or more consecutive amino acids of the sequence of (a) or (b), with the proviso that said subsequence does not fall within amino acid nos. 1102 and 1152 of SEQ ID NO:2.

Preferably, the amino acid sequence in (b) comprises a conservative substitution of at least one amino acid of the amino acid sequence of SEQ ID NO: 2. More preferably, the polypeptide fragment has stem cell differentiation inducing activity, as described above. Useful fragments may exhibit an epitope recognized by polyclonal antibodies raised against the polypeptide having the amino acid sequence shown in SEQ ID NO: 2. for example. A particularly preferred polypeptide is that encoded by the amino acid sequence shown in SEQ ID NO: 2, derived from mouse tissue.

Therefore, also included within the invention are variants and derivatives of the polypeptide described by SEQ ID NO:2 or fragment thereof, whether produced by recombinant means or synthetic means or isolated from naturally occurring sources. For example, peptides having modified amino acids/peptide linkages, and peptides containing non-naturally occurring amino acids and/or cyclic peptides, which may have improved properties such as stability or activity are included. In addition the peptides of the invention may be in the form or a fusion with another protein, for example, tags for the targeted delivery or detection, or purification of the polypeptide (including fragments thereof).

A "variant" in terms of amino acid sequence defines an amino acid sequence that differs by one or more amino acids from another, usually related amino acid sequence. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g.

5 replacement of leucine with isoleucine). Less likely, a variant may have "non-conservative" changes, e.g. replacement of a glycine with a tryptophan.

Similar minor variations may also include amino acid deletions or insertions (i.e. additions), or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing

10 activity (e.g., anti-cancer activity, osteoblast promoting activity, antigenic activity) may be found using computer programs well known in the art.

Variants of the polypeptides of the invention include all forms of mutant variants, for example wherein at least one amino acid is deleted or

15 substituted. Any changes involving substitution of amino acids are preferably neutral or conservative substitutions. Other variants include proteins or polypeptides comprising at least one additional amino acid in the sequence, and/or further comprising an additional amino acid sequence or domain, such as fusion proteins, well known in the art.

20 Further variants of the polypeptides of the invention include those wherein at least one of the amino acids in the sequence is a natural or unnatural analogue. Also, one or more amino acids in the sequence may be chemically modified, e.g. to increase physical stability or to lower susceptibility to enzymic, particularly protease or kinase, activity.

25

Also provided are antibodies that are specifically reactive against the polypeptides of the invention. Methods for producing antibodies are well known in the art. An antibody specific for the polypeptide of the invention can be easily obtained by immunizing an animal with an immunogenic amount of

30 the polypeptide. Therefore, an antibody recognizing the polypeptide of the invention embraces polyclonal antibodies and antiserum which are obtained by immunizing an animal, and which can be confirmed to specifically recognize the polypeptide of the invention by Western blotting, ELISA, immunostaining or other routine procedure known in the art.

It is well known that if a polyclonal antibody can be obtained by sensitization, a monoclonal antibody secreted by a hybridoma may be obtained from the lymphocytes of the sensitized animal (Chapter 6, Antibodies A Laboratory  
5 Manual, Cold Spring Harbor Laboratory Press, 1988). Therefore, monoclonal antibodies recognizing the polypeptide of the invention are also provided. Methods of producing polyclonal and monoclonal antibodies are known to those of skill in the art and described in the scientific and patent literature, see, e.g., Coligan, Current Protocols in Immunology, Wiley/Green, NY (1991);  
10 Stites (eds.) Basic and Clinical Immunology (7th ed.) Lange Medical Publications, Los Altos, CA, and references cited therein (Stites); Goding, Monoclonal Antibodies: Principles and Practice (2nd ed.) Academic Press, New York, NY (1986); and Kohler (1975) Nature 256: 495. Such techniques include selection of antibodies from libraries of recombinant antibodies  
15 displayed in phage or similar on cells. See, Huse (1989) Science 246: 1275 and Ward (1989) Nature 341: 544. Recombinant antibodies can be expressed by transient or stable expression vectors in mammalian cells, as in Norderhaug (1997) J. Immunol. Methods 204: 77-87.

20 In this invention, an antibody also embraces an active fragment thereof. An active fragment means a fragment of an antibody having activity of antigen-antibody reaction. Specifically named, these are active fragments, such as F(ab')<sub>2</sub>, Fab', Fab, and Fv. For example, F(ab')<sub>2</sub> results if the antibody of this invention is digested with pepsin, and Fab results if digested with papain. Fab'  
25 results if F(ab')<sub>2</sub> is reduced with a reagent such as 2-mercaptoethanol and alkylated with monoiodoacetic acid. Fv is a mono active fragment where the variable region of heavy chain and the variable region of light chain are connected with a linker. A chimeric antibody is obtained by conserving these active fragments and substituting the fragments of another animal for the  
30 fragments other than these active fragments. In particular, humanized antibodies are envisioned.

The nucleic acid and polypeptide sequences investigated herein have been found to be differentially expressed in samples obtained from metastatic

cancer cell lines and are predictive of tenascin-W expression in metastatic cancer tissue, as well as in other types of cancer and diseases.

Accordingly, certain aspects of the present invention relate to nucleic acids  
5 differentially expressed in tumour tissue, especially metastatic cancer cell  
lines, polypeptides encoded by such nucleic acids, and antibodies  
immunoreactive with these polypeptides, and preparations of such  
compositions. Moreover, the present invention provides diagnostic and  
therapeutic assays and reagents for detecting and treating disorders involving,  
10 for example, aberrant expression of the subject nucleic acids.

Thus, in a further aspect of the invention, a composition is provided  
comprising an isolated nucleic acid molecule encoding tenascin W or a  
fragment thereof and a pharmaceutically acceptable excipient, diluent or  
15 carrier. The pharmaceutical use of nucleic acids encoding tenascin W has not  
previously been suggested and therefore in this embodiment, the nucleic  
acids of the pharmaceutical compositions are not limited to the nucleic acids  
of the invention. In particular, the composition may comprise an isolated  
nucleic acid having a nucleotide sequence selected from the group consisting  
20 of:

- (a) a nucleotide sequence as set forth in SEQ ID No. 1 or SEQ ID No. 3 (encoding human tenascin.W);
  - (b) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4;
  - 25 (c) a nucleotide sequence with at least 35% identity, preferably at least 40, 50, 60, 70, 80, 90, 95 or 100% identity to any one of the sequences of (a) or (b), preferably (a);
  - (d) a subsequence of at least 10, 15, 20, 25, 30, 40, 50, 75, 100 or more consecutive nucleotides of the sequence of (a),(b) or (c); and
  - 30 (e) a nucleotide sequence complementary to (a),(b), (c), or (d),
- and a pharmaceutically acceptable excipient, diluent or carrier.

In one embodiment, the nucleic acid molecule encodes tenascin-W having the amino acid sequence as set forth in SEQ ID No. 2 or SEQ ID NO. 4 or an



amino acid with at least 30%, preferably at least 50%, 70%, 80%, 90%, 95%, or 100% identity to a sequence corresponding to SEQ ID NO:2 or 4. The nucleic acid molecules are at least 10, preferably at least 15, 20, 30, 50, 75, 100 or more consecutive nucleotides of SEQ ID No. 1 or SEQ ID No.2 or a  
5 sequence complementary thereto.

In one embodiment, the invention provides a composition comprising a nucleotide sequence fragment selected from the group consisting of nucleotides 2380-3171 of SEQ ID No:1 or nucleotides 2371-3162 of SEQ ID  
10 No:3, a complement of nucleotides 2380-3171 of SEQ ID No:1 or of nucleotides 2371-3162 of SEQ ID No:3, and RNA equivalents thereof, which encode an epitope for the binding with an antibody paratope.

In another embodiment, the nucleic acid molecule preferably encodes a  
15 protein having stem cell differentiation inducing activity. Although it is well within the skill of the art to identify polypeptides with an appropriate activity using routine methodology, the isolated nucleic acid molecule preferably encodes the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO:4, most preferably that of SEQ ID NO:4.

20

In yet another embodiment, the nucleic acid molecule has a subsequence that is antisense to SEQ ID NO:1 or SEQ ID NO:3, wherein the nucleic acid molecule may comprise nucleotide residues that are resistant to nuclease degradation.

25

The nucleic acid may be antisense to all or a part of a nucleic acid which hybridizes under stringent conditions to SEQ ID No:1 or SEQ ID No:3, or antisense to a sequence having at least 70% identity with SEQ ID NO:1 or SEQ ID NO:3, that is able to hybridize under low stringency conditions to SEQ  
30 ID NO:1 or SEQ ID NO:3, and which encodes tenascin-W. Low stringency conditions employs around 0.01 x SSC buffer compared to high stringency which employs about a 10 fold greater concentration. Alternatively, the antisense RNA may be antisense to regulatory sequences of the tenascin-W gene, in particular to 5' upstream sequences (promoter region) of the gene.

Similarly, small RNAi oligonucleoties can be designed to inhibit expression of Tenascin-W in a specific manner.

5 The nucleic acids can be RNA or DNA, sense or antisense, and in some embodiments, double stranded (including siRNA) or single stranded. In certain embodiments at least some of the nucleotide residues of the nucleic acids (sense or antisense) may be made resistant to nuclease degradation and these can be selected from residues such as phosphorothioates and/or methylphosphonates.

10

The antisense nucleic acids as hereinbefore described can advantageously be used as pharmaceuticals, preferred pharmaceutical applications being for the manufacture of a medicament for the prophylaxis or treatment of conditions dependent on elevated Tenascin W levels, such as cancer.

15

Thus, the invention provides a method of preventing or treating a condition dependent on Tenascin W, comprising administering to an individual an effective amount of a nucleic acid, as hereinbefore described. Thus, the invention encompasses the use of such nucleic acid molecules as a  
20 pharmaceutical, as well as for the manufacture of a medicament, in particular for the prophylaxis or treatment of cancer or bone pathologies.

25

In yet another aspect, the present invention provides expression vectors capable of replicating in a host cell, comprising one or more vector sequences and a nucleic acid sequence encoding teneurin-W. The construct for use as a pharmaceutical is also provided, as well as its use for the manufacture of a medicament for the prophylaxis or treatment of cancer or the prophylaxis or treatment of bone pathologies.

30

Other embodiments of the invention include nucleic acid constructs capable of replicating in a host cell, comprising (a) at least one nucleic acid sequence portion encoding a tenascin-W protein or polypeptide of the invention (b) antisense nucleic acids as hereinbefore described (or their complement, for example, if expression of the antisense RNA in a cell is foreseen), or (c)

nucleic acids as hereinbefore described and at least one nucleic acid sequence encoding a protein other than tenascin-W (or its homologues), e.g. vector sequence. Such constructs are not naturally occurring sequences. The constructs lack essential sequences of DNA which might permit them to function as vectors but are not naturally occurring as "hybrid" nucleic acids. They may include nucleic acid sequences that function as linkers or restriction sites which include without limitation a transcriptional regulatory sequence operably linked to a nucleotide sequence of the invention so as to render said nucleic acid construct capable of replicating in a host cell. Preferred constructs are synthesised using methods of oligonucleotides synthesis well known to those of skill in the art.

Also provided are vectors comprising a construct as hereinbefore described. Preferred vectors are expression vectors, preferably plasmids or viruses although cloning vectors are also provided for, optionally in the form of plasmids.

The invention provides host cells containing vectors. Preferred host cells are eukaryotic cells, more preferably insect cells or mammalian cells.

Constructs, vectors and transformed host cells of the invention are of use as pharmaceuticals, as well as for the manufacture of a medicament for the prophylaxis or treatment of a condition dependent on tenascin W, such as cancer or bone disorders.

Similarly, in a further aspect of the invention, a composition is provided comprising tenascin W, preferably recombinant tenascin-W, or a fragment thereof and a pharmaceutically acceptable excipient, diluent or carrier. In preferred embodiments, the tenascin-W is a polypeptide having an amino acid sequence selected from the group consisting of:

- (a) an amino acid sequence as set forth in SEQ ID No. 2 or 4;
- (b) an amino acid sequence with at least 35% identity, preferably at least 50%, 70%, 80%, 90%, 95%, or 99% identity to the sequence of (a); and

(c) a subsequence of at least 5, 10, 15, 20, 30, 50, 75, 100 or more consecutive amino acids of the sequence of (a) or (b).

Preferably, the polypeptide has stem cell differentiation inducing activity as described above. More preferably, the polypeptide is encoded by the amino acid sequence shown in SEQ ID NO: 4.

Thus, also provided is the use of tenascin-W for the treatment or prophylactic treatment of any disease or condition requiring increased tenascin-W levels, e.g. thrombosis, wound healing or atherosclerosis, as well as a condition ameliorated by the promotion of osteogenesis, e.g. bone healing, osteoporosis, as well as the use of tenascin-W as a stem cell marker. In yet a further aspect, the tenascin-W protein is used as a pharmaceutical.

The present invention further provides the use of a tenascin-W, e.g. for the manufacture of a medicament, for the prevention or prophylactic treatment of tumourigenesis or the treatment or prophylactic treatment of tumours or cancer. The invention also includes the use of the tenascin-W for the manufacture of a medicament for the treatment or prophylactic treatment of any one or more of bone disease, rheumatism, asthma, allergic diseases, autoimmune diseases, prevention of transplant rejection and any other diseases involving tenascin e.g. thrombosis, cancer, wound healing and arteriosclerosis.

The invention therefore provides pharmaceutical compositions for humans or veterinary compositions for animal use that comprise one or more of the aforementioned active fragments of tenascin-W. The compositions may also include other active or non-active agents. Non-active agents may include a pharmaceutically acceptable excipient, diluent or carrier, but not limited to saline, buffered saline, dextrose and water.

The compositions and medicaments of the invention may therefore be used prophylactically in order to prevent tumours from forming, or they may be used in a curative or partly curative way to treat or contain a pre-existing tumourous

condition: As well as tumours, cancerous or malignant conditions may be prevented or treated with compositions or medicaments of the invention.

In a particular aspect, the present invention provides the use of the nucleic acid or proteins or polypeptides as hereinabove described, for  
5 radioimmunotherapy. Use of radiolabeled antibody is a promising approach to target radiotherapy directly into the tumor. Anti-tenascin-C antibodies are currently tested in phase I and II clinical trials. Patients with malignant gliomas were administrated locoregional radioimmunotherapy (LR-RIT) using  $^{131}\text{I}$ -  
10 labeled anti-tenascin antibody injected directly in the tumor (Riva et al., 1999a). The first results show that LR-RIT can be safely performed, with good results especially in patients with minimal disease. Similar approach was performed with  $^{90}\text{Y}$  (a pure beta emitter)-labeled antibodies Riva et al.,  
15 1999b), with promising results. Potentially more efficient radioimmunotherapies were shown to be possible using other isotopes, like in the case of an  $^{211}\text{At}$ -labeled anti-tenascin antibody (Zhang et al., 2001), without excessive toxicity for the patient. It is as well a useful tool for precise imaging of tumors, since the presence of isotope specifically targeted into the tumor allows sequential scintigraphies of the tumor (Riva et al., 1999a), and  
20 makes possible a direct estimation of the success of the therapy. Similar methodologies can be applied using antibodies specific for tenascin-W.

The tumours or tumour cells of the present invention are preferably those which express tenascin-W in the stroma. In particularly preferred  
25 embodiments the tumours are solid tumours, e.g. mesenchymal tumours such as osteosarcoma, glioblastoma or epithelial cancers such as breast, prostate, lung and colorectal carcinoma.

The invention further provides the use of tenascin-W for the treatment or  
30 prophylactic treatment of a condition ameliorated by the promotion of osteogenesis, e.g. osteoporosis, osteoarthritis, treatment of cartilage and bone pathologies. A protein or polypeptide as hereinabove described may be used to be incorporated into implants including without limitation hip joints, knee joints, or broken bones, to promote osteogenesis.

The invention also provides a method of preventing or prophylactic treatment of tumourigenesis or of treatment or prophylactic treatment of tumours or cancer or of any one or more of rheumatism, asthma, allergic diseases, autoimmune diseases, prevention of transplant rejection or the treatment or prophylactic treatment of any disease involving tenascin-W, e.g., thrombosis; wound healing and atherosclerosis in an individual comprising administering an effective amount of a tenascin-W or a fragment thereof.

- 10 The invention also provides a method of treatment or prophylactic treatment of a condition ameliorated by the promotion of osteogenesis, e.g osteoporosis, osteoarthritis, treatment of cartilage and bone pathologies in an individual comprising administering an effective amount of tenascin-W or a fragment thereof.

15

The determination of an effective dose is well within the capability of those skilled in the art. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in an appropriate animal model. The animal model is also used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

20

A therapeutically effective dose refers to that amount of active agent which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals (e.g.,  $ED_{50}$ , the dose therapeutically effective in 50% of the population; and  $LD_{50}$ , the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio,  $LD_{50}/ED_{50}$ .

25

- 30 Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage varies within this range

depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage may be chosen by the individual physician in view of the patient to be treated. Dosage and administration can be adjusted to provide sufficient levels of the active moiety or to maintain the desired effect.

Additional factors which may be taken into account include the severity of the disease state (e.g. tumour size and location); age, weight and gender of the patient; diet; time and frequency of administration; drug combination(s);

reaction sensitivities; and tolerance/response to therapy. Long acting pharmaceutical compositions can be administered on a daily basis, every 3 to 4 days, every week, or once every two weeks, depending on half-life and clearance rate of the particular formulation.

The present inventors have observed stem cells, especially the periosteum, the bone marrow cell compartment for osteogenesis, expressing tenascin-W and therefore also encompassed by the invention is a method of the invention,

tenascin-W is used as a stem cell marker for cells including without limitation osteogenic precursor cells in the bone marrow. Therefore, also

provided is a method of selecting stem cells or progenitor cells having the ability to differentiate into osteoblasts from other cells, not having this ability.

Stem cells expressing tenascin-W can be detected by an antibody. The antibody recognizing tenascin-W can be detected using secondary antibodies specific for the tenascin-W antibody, which are optionally labelled with a

radiolabel, an enzyme, avidin or biotin, or fluorescent materials (e.g. green fluorescent protein (GFP) or rhodamine), for example. The cells are

characterized by having tenascin-W expression above basal levels and are preferably selected from a mixed population of cells using the fluorescence-activated cell sorter (FACS) (see for example Abe et al., Dev Biol. 1996;

180(2):468-72). The selected cells therefore carry a protein detectable by fluorescence. The sorted cells are useful for the production of biological parts of the body, e.g. bone tissue.

Also provided are antibodies that specifically recognizes tenascin W for use as a pharmaceutical, as well as for the manufacture of a medicament, for the prophylaxis or treatment of cancer (e.g., glioblastoma, prostate, lung, colorectal, osteo- or breast carcinoma), including metastatic cancer, or for the  
5 prophylaxis or treatment of any disease or condition involving tenascin-W, e.g. excessive bone growth. In another aspect of the present invention, an antibody specifically reactive against tenascin-W or a fragment thereof, and the use of an antibody for the manufacture of a medicament for the prophylaxis or treatment of cancer, and the antibody for use as  
10 pharmaceutical is provided.

Antibodies that specifically recognize tenascin-W or a fragment thereof are also provided, in particular antibodies that recognise the above mentioned epitope.  
15

Methods for detecting tenascin-W embrace, for example, the use of an antibody as referred to above, optionally with the use of an enzyme reaction. The antibody recognizing tenascin-W can be detected using secondary antibodies specific for the tenascin-W antibody, which are optionally labelled  
20 with a radiolabel, an enzyme, avidin or biotin, or fluorescent materials (FITC or rhodamine), for example.

Also encompassed by the invention is the use of an antibody that specifically recognizes tenascin-W for the manufacture of a medicament, in particular a  
25 medicament for the prophylaxis or treatment of cancer, the prophylaxis or treatment of bone disease, or as a pharmaceutical. In particularly encompassed by the invention is the use of an antibody that specifically recognizes tenascin-W for the diagnosis of tumour, especially metastatic tumour.

30 In a further embodiment, the present invention provides a method for identifying agents for the prevention or the prophylactic treatment of tumourigenesis or the treatment or prophylactic treatment of tumours or cancer, or the treatment or prophylactic treatment of any disease or condition



involving tenascin-W, e.g. a condition ameliorated by the promotion (or inhibition) of osteogenesis, comprising contacting a test compound with a tenascin-W expressing sample and then measuring a change in one or more of (a) cell proliferation, e.g. cell progression entering S-phase of the cell cycle; (b) DNA synthesis; (c) cell adhesion; (d) cell spreading; (e) focal adhesion and actin stress fibre formation on fibronectin; (f) cell binding to extracellular matrix (ECM), relative to when said test compound is absent.

Cells may be encouraged to proliferate by the addition of tenascin-W to the cell culture, preferably by coating the solid substrate therewith. A substrate can be any surface that promotes cell adhesion. The solid substrate may also be coated by other ECM which include without limitation fibronectin, collagen, etc. The cell cultures are preferably grown on a solid substrate or in a liquid medium. A first measurement of one or more of (a) to (f) may be made prior to contacting the cells with a test substance. A second measurement may be made thereafter. A multiplicity of further measurements may be made over a period of hours or days after contact of the cells with the test compound. In this way a time course of the cellular response(s) may be determined and analysed.

20

In one preferred embodiment of the present invention, the presence of tenascin-W in the liquid medium is measured relative to when a test compound is absent. An increase in the level of tenascin-W present in the medium relative to when said test agent is absent correlates to an agent effective in the promotion of osteogenesis, for example. A decrease in the level of tenascin-W present in the medium relative to when said test agent is absent correlates to an anti-proliferative or anti-tumour agent, or an agent effective in inhibiting osteogenesis or osteoblast formation.

30

In preferred aspects one or more of the following conditions arising after contacting cells with a test compound is indicative of an anti-proliferative or anti-tumour agent, or an inhibitor of osteoblast formation:

- 5
- (a) a reduction in cell proliferation; or a decrease in the proportion of cells entering S-phase of the cell cycle;
  - (b) a reduction in DNA synthesis;
  - (c) an increase in cell adhesion;
  - (d) an increase in cell spreading;
  - (e) an increase in focal adhesion and actin stress fibre formation on fibronectin; and
  - (f) an increase in the binding of cells to ECM, preferably fibronectin;

10

In other preferred aspects one or more of the following conditions arising after contacting cells with a test compound may indicate an osteogenesis promoting agent:

15

- (a) an increase in cell proliferation; or an increase in the proportion of cells entering S-phase of the cell cycle;
- (b) an increase in DNA synthesis;
- (c) and (d) an increase in expression of bone-specific markers such as alkaline phosphatase activity, calcification or any others known in the art (e.g., Raouf and Seth, 2002, Bone 30: 463-71).

20

Actin stress fibre formation may be assayed according to the Actin Assembly Assay described in Bloom, L *et al* (1999) Mol Biol Cell 10: 1521-1536.

25

Adhesion assays may be performed according to the method described in Bloom, L *et al* (1999).

30

In other embodiments, the method of the invention may further comprise control cells grown in the absence of test substance and (a), (b), (c), (d), (e), and/or (f) are measured in both control and test cultures. The test measurements can thereby be normalised with respect to the control.

The screening method further provides an essentially cell-free system for the identification of potential anti-tumour or tumour preventing agents or for an agent inhibiting osteogenesis. This method relies on the ability of a potential

anti-tumour agent to prevent, inhibit or disrupt the binding between an ECM and tenascin-W. The nature of any disruption of the ECM and tenascin-W binding may be determined by performing a binding assay for ECM and tenascin (see e.g. example 10). For example, calorimetric methods may be  
5 used or measurement of labelled reagents.

Alternatively, a method is provided for identifying modulators of tenascin W function, comprising: (a) contacting a test compound with tenascin W and/or alpha8 beta1 integrin, and (b) measuring the binding of the test compound to  
10 tenascin-W and/or alpha8 beta1 integrin, or (c) measuring a disruption of binding of tenascin-W to alpha8 beta1 integrin, relative to when the test compound is absent. A decrease in binding of tenascin-W to alpha8 beta1 integrin is indicative of an inhibitor of this interaction, and increased binding could indicate that the test compound activates the alpha8 beta1 integrin,  
15 thereby increasing the interaction between tenascin-W and alpha8 beta1.

The relative amounts or concentrations of reagents and test substance may be varied, thereby permitting calculation of inhibition constants and other parameters, e.g. binding affinities. The optimisation of assay conditions will  
20 be well within the realm of one of ordinary skill in the art. The system may further comprise a control without test substance and the binding is measured in the control, thereby permitting corresponding measurements in the test system to be normalised relative to the control.

25 Where one component of the assay (screening) systems of the invention is coupled to a solid particle or substrate, then one or more of the other components not so coupled may be labelled. Examples of labels include radiolabels e.g.  $^{14}\text{C}$  or  $^3\text{H}$ , dyes, metal sols, enzymes or biotin/avidin. By attaching such labels to "free" components in the system any binding assay  
30 may be carried out in solution in accordance with procedures well known in the art. After allowing the components to react solid phase particles can be separated from solution, e.g. by filtration or sedimentation (including centrifugation). In some embodiments immunoprecipitation may be used to separate bound and free labelled components. Usually, an antibody may be

employed to bring an unlabelled component out of solution (whether or not this component has bound to another labelled component or not). After separation, the label present in solution (free) and the label present in or on the solid phase (bound) may be measured. Standard analyses of such bound and free data, e.g. Scatchard plots and the determination of affinity and inhibition constants for binding are well known to the person of ordinary skill in the art.

Where the solid phase is not particulate, e.g. in the form of a surface, such as a microtiter plate well, then binding assays measuring bound and free label may be performed but this will normally involve the removal of liquid phase from the wells after binding reactions have occurred. Advantageously, this assay format may dispense with the need for providing specifically labelled reaction components. Instead, labelled antibodies may be used to measure the binding of previously free reaction components to solid phase components.

In some embodiments the tenascin-W molecule, variant or fragment thereof may be attached directly to a solid phase. In preferred immunoassay embodiments of this type, tenascin-W bound to an ECM is measured using an antibody reactive against tenascin-W.

Immunological binding assays are known in the art. For a review, see *Methods in Cell Biology Vol. 37: Antibodies in Cell Biology*, Asai, (Ed.) Academic Press, Inc. New York (1993).

A label may be any detectable composition whereby the detection can be spectroscopic, photochemical, biochemical, immunochemical, physical or chemical. For example, useful labels can include  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , fluorescent dyes (e.g. FITC, rhodamine and lanthanide phosphors), electron-dense reagents, enzymes, e.g. as commonly used in ELISA (e.g. horseradish peroxidase, beta-galactosidase, luciferase and alkaline phosphatase), biotin, dioxigenin, or haptens and proteins for which antisera or monoclonal antibodies are available. The label may be directly incorporated into a target

compound to be detected, or it may be attached to a probe or antibody which binds to the target.

Throughout the assays of the invention, incubation and/or washing steps may be required after each application of reagent or incubation of combinations of reagents. Incubation steps may vary from about 5 minutes to several hours, perhaps from about 30 minutes to about 6 hours. However, the incubation time usually depends upon the assay format, analyte, volume of solution, concentrations, and so forth. Usually, the assays should be carried out at ambient temperature, although they may be conducted at temperatures; in the range 10°C to 40°C, for example.

A particularly preferred assay format is an enzyme-linked immunosorbent assay (ELISA).

All of the aforementioned methods of screening of the invention are equally applicable to the screening of substances for biological activity and potential agents for any other disease or condition involving tenascin-W, e.g. wound healing or treatment of arteriosclerosis.

Also included within the scope of the present invention are anti-tumourigenic, anti-tumour, anti-metastatic, (anti-)osteogenic, wound healing or anti-arteriosclerosis substances or substances for the treatment or prophylactic treatment of any disease or condition involving tenascin-W identified by any of the screening methods of the invention. These substances may be proteins, polypeptides or small organic molecules (drugs). The invention therefore includes pharmaceutical compositions for preventing or treating tumours, metastasis, or bone pathologies and comprising one or more of the substances identified by a method of the invention. For example, inhibitors of tenascin-W expression or activity are considered potential anti-cancer agents, whereas tenascin W or agonists thereof are considered agents effective in promoting osteogenesis, which can be used in vivo or ex vivo.

Thus, the present invention provides a novel mammalian member of the tenascin family and uses thereof. It permits the identification of agents effective against conditions dependent on tenascin-W, in particular anti-cancer agents or agents that promote osteogenesis, by performance of any of the methods of screening described herein. Preferred anti-cancer agents are those which inhibit proliferation of the cancer cells and which may be general anti-proliferative agents.

The invention includes all nucleic acid molecules and proteins and polypeptides as hereinabove described, as well as agents identified by performing the methods, and the use of these agents as pharmaceuticals, particularly as medicaments for the prophylaxis or treatment of cancer and other conditions dependent on tenascin W.

Thus, in a further aspect the invention provides for the use of tenascin-W and of an agent identified by a screening method of the invention as a pharmaceutical.

The invention further provides tenascin-W or an agent identified by a screening method of the invention, for the manufacture of a medicament for the prophylaxis or treatment of a condition dependent on tenascin-W, for use to treat cancer or bone diseases or an immunological defect.

The invention provides a method of preventing or treating a condition dependent on tenascin-W comprising administering to an individual an effective amount of a construct, vector, host cell or antibody described above.

The invention also provides a method of inhibiting a condition dependent on tenascin-W comprising administering an effective amount of the modulator identified by a screening method of the invention described above, for the treatment of cancer or bone disease or an immunological defect.

Also provided by the invention are the nucleic acid molecules, the proteins, and the agents referred to above in a pharmaceutical composition, possibly in

the presence of suitable excipients known to one of ordinary skill in the art.

The compositions may be administered in the form of any suitable composition as detailed below by any suitable method of administration within the knowledge of a skilled man. The preferred route of administration is

5 parenterally. In parenteral administration, the compositions of this invention will be formulated in a unit dosage injectable form such as a solution, suspension or emulsion, in association with a pharmaceutically acceptable excipient. Such excipients are inherently nontoxic and nontherapeutic.

10 Examples of such excipients are saline, Ringer's solution, dextrose solution and Hank's solution. Nonaqueous excipients such as fixed oils and ethyl oleate may also be used. A preferred excipient is 5% dextrose in saline. The excipient may contain minor amounts of additives such as substances that enhance isotonicity and chemical stability, including buffers and preservatives.

15 Any protein is administered at a concentration that is therapeutically effective to prevent organ rejection, GVHD, allergy and autoimmune diseases. The dosage and mode of administration will depend on the individual. Generally, the compositions are administered so that the functional protein is given at a dose between 1 pg/kg and 10 mg/kg, more preferably between 10 ug/kg and  
20 5 mg/kg, most preferably between 0.1 and 2 mg/kg. Preferably, it is given as a bolus dose. Continuous short time infusion (during 30 minutes) may also be used. The compositions according to the invention may be infused at a dose between 5 and 20 µg/kg/minute, more preferably between 7 and 15 µg/kg/minute.

25

According to a specific case, the "therapeutically effective amount" of a composition needed should be determined as being the amount sufficient to cure the patient in need of treatment or at least to partially arrest the disease and its complications. Amounts effective for such use will depend on the  
30 severity of the disease and the general state of the patient's health. Single or multiple administrations may be required depending on the dosage and frequency as required and tolerated by the patient.

The present invention also provides a method of diagnosing or prognosing cancer, or any other condition dependent on elevated tenascin W levels, comprising, (a) analysing a sample obtained from an individual for the presence of tenascin-W; and (b) correlating the presence of tenascin-W with  
5 an unfavourable prognosis or diagnosis.

The methods of the present invention will typically involve the determination of the presence, level, or activity of tenascin-W in a cell or tissue sample, which sample will often be obtained from a human, but one can also readily  
10 understand that samples tested by the present method can be obtained from agriculturally important mammals, such as cattle, horses, sheep, etc., or other animals of veterinary interest, such as cats and dogs. The assay may be carried out on any cell or tissue sample, such as somatic tissues, germine  
15 tissues, or cancerous tissues, as well as on samples from body fluids, such as pleural fluid, blood, serum, plasma and urine. The method may also further comprise propagating cells in the sample in cell culture.

A "sample" is the material being analyzed which is usually, but not necessarily, subjected to pretreatment to provide the tenascin-W in assayable  
20 form. This would for example, entail forming a cell extract, methods for which are known in the art (for example, see Scopes, Protein Purification: Principles and Practice, Second Edition (Springer-Verlag, N.Y., 1987)).

In the broader aspects of the invention, there is no limitation on the collection  
25 and handling of samples as long as consistency is maintained. The sample is obtained by methods known in the art, such as, biopsies, surgical resections, smears, or the like. Optionally, cells obtained in a sample may be propagated in cell culture.

30 Consistency of measurement of tenascin-W or tenascin-W activity in clinical samples can be ensured by using a variety of techniques. For example, to control for the quality of each tissue extract, another enzymatic activity, such as alkaline phosphatase, can serve as an internal control. In addition, an internal standard can be measured concurrently with tenascin-W in the



sample as a control for assay conditions. Thus, the analyzing step can comprise detecting a control protein in the sample, optionally normalizing the value obtained for tenascin-W with a signal obtained with the control protein.

- 5 The presence of tenascin-W in the sample can be determined by detecting the tenascin-W protein using methods known in the art. In this invention, there are no limitations on the type of assay used to measure tenascin-W or tenascin-W activity. For example, tenascin-W can be detected by immunoassays using antibodies specific for tenascin-W. The antibody can be used, for example, in
- 10 Western blots of two dimensional gels where the protein is identified by enzyme linked immunoassay or in dot blot (Antibody Sandwich) assays of total cellular protein, or partially purified protein.

- Methods for sample concentration and protein purification are described in the
- 15 literature (see Scopes, 1987). For example, if desired, the tenascin-W present in the cell extract can be concentrated, by precipitating with ammonium sulfate or by passing the extract through a commercially available protein concentration filter, e.g., an Amicon or Millipore, ultrafiltration filter. The extract can be applied to a suitable purification matrix, such as an anion or a cation
- 20 exchange resin, or a gel filtration matrix, or subjected to preparative gel electrophoresis. In such cases, the tenascin-W and protein yield after each purification step needs to be considered in determining the amount of tenascin-W in a sample.

- 25 Tenascin-W may be detected using an antibody specific for tenascin-W, and a control assay can be carried out using an antibody specific for another tenascin molecule. Optionally, the method may further comprise correlating in an increase in tenascin-W in the sample relative to healthy tissue. For example, tenascin-W can be detected using an antibody specific for tenascin-
- 30 W expressed in tumour tissue and compared to antibody binding to any tenascin-W expressed (or non-specific reaction) in healthy tissue.

The sample is preferably a tissue sample mounted onto a solid surface for histochemical analysis. The presence of detectable, accessible tenascin-W

indicates that tenascin-W is accessible to cells for binding. This leads to a unfavourable diagnosis or prognosis. If, on the other hand, the antibody does not react with tenascin-W in the tissue section, then there is an expectation that tenascin-W is not present. This leads to a favourable diagnosis or prognosis.

The present inventors have found that tenascin-W is specifically expressed in solid tumours, in particular metastatic tumour tissue or stroma thereof. The presence of tenascin-W therefore indicates a cancerous condition, in particular the presence of metastatic tumour tissue, whereas the absence of tenascin-W indicates healthy tissue or non-metastatic tumour tissue. Tenascin-W was identified in developing mouse tissues by western blotting. High expression of tenascin-W was found in the metastatic tumours of ras-transgenic mice, but not in the myc- or neuT-transformed non-metastatic tumours. The presence of tenascin-W (170 kD) is indicative of unfavourable diagnosis.

In a further embodiment, the diagnostic methods of the invention further comprises analysing the sample for the presence of alpha 8 integrin, the presence of alpha 8 integrin correlating with an unfavourable prognosis or diagnosis. This can easily be achieved, for example, using an antibody as described in detail in Example 8 below.

In a preferred embodiment, the invention provides kits suitable for use in the diagnostic or prognostic methods of the invention. Such kits comprise reagents useful for carrying out these methods, for example, antibodies from one or more species specific for tenascin-W and alpha 8 beta1 integrin. Secondary antibodies that recognise either or both such primary anti-fibronectin antibodies can also be included for the purpose of recognition and detection of primary antibody binding to a sample. Such secondary antibodies can be labelled for detection e.g. with fluorophores, enzymes, radioactive labels or otherwise. Other detection labels will occur to those skilled in the art. Alternatively, the primary anti-tenascin-W antibodies can be labelled for direct detection.

The invention is further described below, for the purpose of illustration only, in the following examples.

**5 Example 1: Cloning of mouse tenascin-W**

Mouse tenascin-W was cloned from a cDNA library of 19d whole mouse embryos (DupLEX-A DLM-1110; OriGene Inc.). In a first step the following PCR primers derived from a sequence from chromosome 1, similar to Tenascin-R (Accession number AL049689) were used for nested PCR reactions with the  
10 Expand High Fidelity PCR System (Roche) using the mouse cDNA library as template. The first reaction was performed with the primer set 5'-TAGCAGCCCACAGCATCTACTTGCC-3' (SEQ ID NO:5) / 5'-ATTGCTGTTCTGCTGAACCTGACTGCA -3' (SEQ ID NO:6) and the second reaction with 5'-ATGGATCCAGAAATTGACGGCCCCAAAACCTAG-3' (SEQ  
15 ID NO:7) / 5'-ATAAGCTTGTGGAGAGGGTGGTGGATACATTTC-3' (SEQ ID NO:8). The primer set included a BamHI and a HindIII restriction site, respectively, to allow directed cloning into the bacterial expression vector pQE30 (Qiagen), employing a C-terminal His-tag for the purification of the recombinant proteins.

20

The mouse proteins (tenascin W polypeptide fragments obtained as a result of the above procedure) were expressed in *E. coli* and purified by affinity chromatography to a Ni-NTA matrix (Qiagen) according to the matrix supplier's manual. The protein was purified under native conditions and was  
25 eluted with 250mM imidazole.

Full length tenascin-W was cloned by the use of mouse tenascin-W specific primers derived from the above mouse tenascin-W cDNA and primers matching the vector of the same 19d whole mouse embryo cDNA library used  
30 before. To obtain the complete 5' sequences, the following PCR reactions using the above cDNA as template were performed: The first PCR reaction was performed using the primer pair 5'-AGGAGATGGTGGCTGTATTTTCGG-3' (SEQ ID NO:9) / 5'-AGCCTCTTGCTGAGTGGAGATGCC-3' (SEQ ID NO:10) followed by a second PCR reaction with the primer set 5'-

TAGAATTCGGTCACCTGATTGGTCACTAGG-3' (SEQ ID NO:11) / 5'-TTATGATGTGCCAGATTATGCC-3' (SEQ ID NO:12). To complete the 3' part of the tenascin-W cDNA the following PCR reactions were performed: In the first reaction the primer pair 5'-CTCAAATTGATGGCTACATTTTGACC-3' (SEQ ID NO:13) / 5'-AAGCCGACAACCTTGATTGGAGAC-3' (SEQ ID NO:14) was used followed by the primer pair 5'-TACCAGTTCCCAAATGGCACCG-3' (SEQ ID NO:15) / 5'-AAACCTCTGGCGAAGAAGTCC-3' (SEQ ID NO:16). In each case the longest products were cloned. These overlapping tenascin-W cDNA clones were assembled into one full length mouse tenascin-W cDNA and cloned into the expression vector pCEP/Pu (see Kohfeldt et al. (1997). FEBS Lett. 414:557-61). At the 3'end of the tenascin-W cDNA a 6xHis-tag was inserted in front of the stop codon to allow the purification of full length mouse tenascin-W protein expressed in mammalian cell culture.

- 15 The recombinant mouse tenascin-W protein comprises three C-terminal fibronectin type III repeats in the region defined by amino acids 794-1057 of the complete amino acid sequence of mouse tenascin-W, encoded by nucleotides 2380 – 3171 of the tenascin-W nucleotide sequence.

## 20 **Example 2: Characterization of mouse Tenascin-W**

- The full length cDNA of mouse tenascin-W was cloned as described in example 1. The cDNA sequence encodes a typical member of the tenascin protein family and harbors from the N-terminus to the C-terminus of the protein the following structural domains: signal peptide for secretion, N-terminal domain for dimerisation of two tenascin-W trimers that are assembled by heptad repeats. This results in a disulfide-linked hexameric protein complex where each subunit contains three and a half EGF-like repeats, nine fibronectin type III repeats, and a fibrinogen-like C-terminal globular domain.

- 30 The full length tenascin-W cDNA was transfected into HEK 293 cells using the transfection reagent fugene (Roche). Transfected cells were selected with puromycin and the medium containing the secreted tenascin-W protein was collected and the protein was purified by sequential chromatography over a gelatin-agarose column (Sigma) to remove any contaminating fibronectin in

the preparation and by adsorption to a Ni-NTA matrix (Qiagen). The tenascin-W was eluted from the nickel column by 250mM imidazole.

The recombinant protein was also analyzed by SDS-PAGE (sodium dodecyl sulfate - polyacrylamide gel electrophoresis) on 6% polyacrylamide gels, under reducing and non-reducing conditions as well as by electron microscopy after rotary shadowing using the same procedure as described for tenascin-C (Chiquet-Ehrismann, R. et al. (1988) Cell 53, 383 – 390).

Tenascin-W showed a similarly slow migration as the hexameric tenascin-C protein. Electronmicrographs of tenascin-W after rotary shadowing indeed revealed hexameric molecules with six subunits of about 50nm length radiating from a central globular domain.

### **Example 3: Cloning of human tenascin-W**

Human tenascin-W was cloned from cDNA made from mRNA isolated from the osteosarcoma cell line Saos-2 (ATCC; HTB 85) essentially as described in Example 1 using the same PCR primers. The human protein was expressed and purified by affinity chromatography to a Ni-NTA matrix (Qiagen) according to the matrix supplier's manual. The protein was purified under native conditions and was eluted with 250mM imidazole.

The recombinant protein comprises the three C-terminal fibronectin type III repeats in the region defined by amino acids 791-1054 of the complete amino acid sequence of human tenascin-W encoded by nucleotides 2371 – 3163 of the tenascin-W nucleotide sequence of the database entry AL049689), respectively.

Full length tenascin-W is cloned by the use of human tenascin-W specific primers derived from the above human tenascin-W cDNA and human genomic sequences 5' to the ATG start codon of the cDNA sequence entry AL049689 using cDNA made from mRNA isolated from osteosarcoma cell line Saos-2 (ATCC; HTB 85) as the template. The following primers are used for three sets of nested PCRs:

hTNW1: 5'CATCCTGGAGGGTCTGCTCC3' (SEQ ID NO:17)



recombinant tenascin-W as well as with endogenous tenascin-W in tissue extracts of mouse organs, as demonstrated by Western blotting. In both cases, tenascin-W was identified as a 170kDa molecular weight species.

- 5 The anti-tenascin-W antiserum was used to investigate tenascin-W expression during normal mouse development by immunohistochemistry. For immunohistochemistry, tissues were fixed in ice-cold 4% paraformaldehyde in phosphate-buffered saline (PBS) overnight, washed with PBS and cryoprotected with 25% sucrose in PBS overnight at 4°C. The tissues were  
10 embedded in OCT (Optimal Cutting Temperature) mounting medium (Cat. No. 27050 OCT Compound by Ted Pella Inc., CA) and sections of 12-16µm were cut and collected onto glass slides. The sections were air-dried for 2 hours before staining with anti-tenascin-W antiserum followed by a fluorescently labelled secondary antibody.

15

- Tenascin-W first appears at embryonic day 11.5 (E11.5) in the maxillary process. Between E14.5 and E16.5, tenascin-W and tenascin-C expression overlaps in developing connective tissue (palate and mandible) in the face  
20 and jaw. Furthermore tenascin-W is found in the extracellular matrix (ECM) of smooth muscle, mesothelia and bone. In the adult mouse tenascin-W is found in a subset of the tenascin-C-positive ECM of the aortic valve and the limbus. In these locations its expression coincides with the stem cell compartment of the respective tissue. Tenascin-W is also expressed in the periosteum, the  
25 kidney and the digestive tract in a subset of tenascin-C-positive regions, but not in the brain.

#### **Example 5: Monoclonal antibodies against human tenascin-W**

- 30 The bacterially expressed recombinant fragment of human tenascin-W as described above in Example 3 was used to raise monoclonal antibodies against human tenascin-W using standard procedures. The monoclonal antibodies reacted specifically with human tenascin-W having better binding

than relying on the crossreactivity of the anti-mouse tenascin-W for human tenascin-W. The monoclonal antibodies are particularly useful to stain human tissues.

**5 Example 6: Tenascin-W expression in tumour cells**

Tenascin-W expression in tumour cells was tested and compared with the known results for tenascin-C which has been found to be highly expressed in tumour tissues (Chiquet-Ehrismann, R. (1993) *Sem. Cancer Biol.* 4, 301-310).

10 Mouse mammary tumours develop readily in transgenic mice expressing oncogenes under the control of mammary gland-specific promoters. Overexpression of c-myc results in the growth of non-metastatic tumours whereas overexpression of Ha-ras leads to the development of metastatic tumours (Li, F. et al. (1994) *Int. J. Cancer* 59, 560-568).

15 In this Example, the antisera described in Example 4 were used to detect tenascin-W in mouse mammary tumours as described for tenascin-Y by Mayes, C. et al. (1996). High expression of tenascin-W (about 170 kDa) was found in the tumours of ras-transgenic mice (metastatic), but not in the myc- or neuT-transformed non-metastatic tumours, whereas tenascin-C was over-  
20 expressed in both types of tumours.

As a control, expression of tenascin-W was examined in healthy tissue, using blood serum, for example. The content of tenascin-W in serum is analyzed by Western blotting. For improved sensitivity a Sandwich ELISA test as  
25 described previously for tenascin-C (Schenk et al. 1995. *Int. J. Cancer* 61:443-449) can be used. Briefly, 96-well plates are coated with either polyclonal or monoclonal anti-tenascin-W antibodies. The serum samples are applied, the wells washed and the bound tenascin-W is detected by either a polyclonal or a monoclonal anti-tenascin-W antibody followed by an  
30 appropriate peroxidase-labeled secondary antibody. No expression of Tenascin-W was found in blood serum from wild-type mice. In contrast, healthy kidney, heart valve and periosteum was found to express Tenascin-W.



Transgenic mice overexpressing neuT develop non-metastasizing mammary tumours, whereas in transgenic mice overexpressing neuT together with EphB4 receptor tyrosine kinase the tumours are metastatic (Munarini, N. et al. (2002) Cell Sci. 115, 25-37). Using this model system we again found high expression of tenascin-W in metastatic tumours, but not in non-metastatic ones. These expression patterns were confirmed by SDS-PAGE (SDS-polyacrylamide gel electrophoresis), by fractionating tumour extracts, blotting on polyvinylidene difluoride membranes, and analyzing the extract using anti-tenascin-W antisera.

#### **Example 7: Adhesion assay**

The purified tenascin-W was used for cell adhesion studies of MDA-MB435 mammary carcinoma cells (ATTC; HTB-129), C2C12 mouse skeletal myoblasts (ATTC; CRL-1772, T98G glioblastoma cells (ATTC; CRL-1690) and NIH-3T3 fibroblasts (ATTC; CRL-1658). In brief, 60-well microtiter plates (Nunc) were coated with 2-100 µg/ml tenascin-W for 1h at 37°C. The non-coated plastic surface was blocked with 1% heat-inactivated BSA in PBS. Cells were trypsinised, trypsin was blocked with 100µg/ml soybean trypsin inhibitor in PBS and, cells were resuspended in serum-free medium and counted. 200-500 cells per well were plated for the indicated time points, fixed by addition of glutaraldehyde (2% final concentration) for 15 minutes and stained with 0.1% crystalviolet in 20% methanol for 30 minutes. Cells were observed under a microscope (Nikon diaphot).

Most cells adhered to tenascin-W coated at 2-100 µg, whereas cell adhesion to tenascin-C was minimal.

We compared the morphology and actin cytoskeleton of C2C12 mouse skeletal myoblasts and T98G glioblastoma plated on tenascin-W to cells plated on fibronectin or tenascin-C. The shape of the cells on tenascin-W was very different from the cells on fibronectin, which became particularly evident after F-actin staining with phalloidin. The cells on fibronectin were well spread containing stress fibers, whereas the cells on tenascin-W had many actin-rich processes but no stress fibers and the cell bodies remained relatively round.

**Example 8: Identification of a cellular tenascin-W receptor**

To determine the cellular receptor(s) responsible for cell adhesion to tenascin-W, we tested the effect of integrin function-blocking antibodies on adhesion of T98G glioblastoma cells on tenascin-W. Antibodies to  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 3$ ,  $\alpha 4$ ,  $\alpha 5$ ,  $\alpha 6$  and  $\alpha V$  were unable to inhibit adhesion of T98G cells to tenascin-W.

Nevertheless, this adhesion was  $\beta 1$  integrin-dependent since 10  $\mu\text{g/ml}$  of the anti- $\beta 1$  integrin blocking antibody P4C10 (Sigma) was able to completely inhibit adhesion to tenascin-W.

IDG tripeptide motifs have been reported to be the recognition sequence for  $\alpha 9 \beta 1$  integrin (Yokosaki et al., 1998). Since mouse tenascin-W contains three IDG motifs we investigated whether  $\alpha 9$  integrin could be the receptor for tenascin-W. We plated SW480 colon carcinoma cells transfected either with an empty vector or with the vector containing the cDNA for  $\alpha 9$  integrin (Yokosaki et al. J Biol Chem. 1998 Sep 27;273(39):24144-50) on tenascin-W coated wells. However, the  $\alpha 9$  transfected SW480 cells failed to adhere to tenascin-W whereas they adhered well to fibronectin and collagen.

Integrin  $\alpha 8$  is expressed in developing rib bone, in kidney and in smooth muscle from the gastro-intestinal tract (Denda et al. Biochemistry. 1998 Apr 21;37(16):5464-74). Since this expression pattern coincides with the presence of tenascin-W it seemed that integrin  $\alpha 8$  was a good candidate receptor for tenascin-W as well. We tested this hypothesis by using the leukemia cell line K562, transfected with  $\alpha 8$  integrin (Denda et al. Biochemistry. 1998 Apr 21;37(16):5464-74). Transfected K562 cells could indeed adhere to tenascin-W and the mock-transfected control cells did not. Therefore,  $\alpha 8 \beta 1$  integrin is a receptor for tenascin-W.

**Example 9: DNA replication and proliferation assay**

96-well plates (Falcon) are coated as described above. Cells are serum starved overnight and trypsinised.  $10^4$  cells are transferred onto the coated

plates in the presence of 1 % serum or 40nM PDGF BB (Platelet-derived growth factor BB). 14h later cells are labelled with radioactive  $^3\text{H}$ -thymidine (0.5 $\mu\text{Ci}$ /well) for 4h at 37°C, incorporated  $^3\text{H}$ -thymidine precipitated with 10% TCA and determined with a Beckman scintillation counter after cell lysis in

5 0.3N NaOH, 2% SDS. Alternatively, incorporation of BrdU is measured or cells numbers are counted over a growth period of several days of cells plated on different substrates. Cancer cells grown on tenascin-W show an increased growth rate over cells plated on fibronectin, as established by counting cells or an increase in radioactive  $^3\text{H}$ -thymidine or BrdU incorporation into cellular

10 RNA.

#### **Example 10: In vitro binding assay (ELISA)**

96-well ELISA plates are coated with the appropriate ECM proteins (e.g. fibronectin or tenascin-W) for 1h at 37°C, blocked with 1% milkpowder, 0.05%

15 Tween-20 in PBS. ECM proteins (tenascin-W or fibronectin) are added in blocking solution. In this assay, an interaction between tenascin-W and fibronectin can be demonstrated. Bound proteins are detected by immune reaction with a peroxidase-coupled secondary antibody followed by

20 colour reaction with 21mg/ml citric acid 1-hydrate, 34mg/ml  $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ , 0.4mg/ml phenylenediamine, 1 $\mu\text{l}$   $\text{H}_2\text{O}_2$ , which can be stopped with 4M sulphuric acid. The absorbance was read at 590nm.

#### **Example 11: Immunofluorescence microscopy**

25  $10^4$  cells are transferred onto 4-well Cellstar plastic plates (Greiner) that are coated with ECM proteins essentially as described above. Cells are fixed with 4% paraformaldehyde, 50mM phosphate buffer, 5mM EDTA in PBS for 15 minutes, blocked with 3% BSA, 0.5% Tween-20 in PBS and incubated with primary and secondary antibodies in blocking solution. Slides are embedded

30 in 10.5% Mowiol containing 2.5% DABCO as antifade agent. Cells are analysed by microscopy. This method is particularly useful for the detection of tenascin-W or any other protein produced by cells in culture to which an antibody is available and can be used to analyze substances that affect the synthesis or deposition of the respective antigens.

As is apparent to one of ordinary skill in the art, variations in the above-described methods can be introduced with ease to attain the same objective.

- 5 Various incubating conditions, labels, apparatus and materials can be chosen according to individual preference. All publications referred to herein are incorporated by reference in their entirety as if each were referred to individually.

**CLAIMS:**

1. An isolated nucleic acid molecule having:

- 5           (a)       a nucleotide sequence as set forth in SEQ ID NO: 1;  
          (b)       a nucleotide sequence encoding a polypeptide having the  
                  amino acid sequence shown in SEQ ID NO: 2;  
          (c)       a nucleotide sequence with at least 85% identity to the  
                  sequence of (a) or (b);  
10          (d)       a subsequence of a least 50 consecutive nucleotides of a  
                  sequence of (a), (b) or (c), with the proviso that said  
                  subsequence does not fall between nucleotide 1027 and  
                  nucleotide 1076 of SEQ ID NO: 1; or  
          (e)       a nucleotide sequence complementary to any of the  
                  nucleotide sequences in (a),(b), (c) or (d).

15           2. The nucleic acid of claim 1 having said nucleotide sequence with  
                  least 85% identity to the sequence of (a) or (b).

20           3. The nucleic acid of claim 1 encoding a polypeptide having the amino  
                  acid sequence shown in SEQ ID NO: 2.

25           4. The nucleic acid of claim 1, having a subsequence of a least 50  
                  consecutive nucleotides, which is antisense to SEQ ID NO:1 or a  
                  sequence having at least 85% identity thereto.

          5. A nucleic acid vector comprising the nucleic acid molecule of any one  
                  of claims 1 to 4.

30           6. A host cell comprising the vector of claim 5.

          7. An isolated polypeptide having:

- (a)       an amino acid sequence as set forth in SEQ ID No. 2;  
          (b)       an amino acid sequence with at least 85% identity to the  
                  sequence of (a); or

- (c) a subsequence of at least 30 consecutive amino acids of the sequence of (a) or (b), with the proviso that said subsequence does not fall within amino acid nos.1102 and 1152 of SEQ ID NO:2.

5

8. The polypeptide of claim 7, wherein said amino acid sequence in (b) comprises a conservative substitution of at least one amino acid in said amino acid sequence of SEQ ID: NO: 2.

10

9. The polypeptide of claim 7 or 8, wherein said polypeptide has stem cell differentiation inducing activity.

10. The polypeptide of claim 7, said polypeptide having the amino acid sequence shown in SEQ ID NO: 2.

15

11. An antibody that specifically recognizes the polypeptide having the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4.

20

12. An antibody that specifically recognizes tenascin W for use as a pharmaceutical.

25

13. The antibody of claim 12, wherein said antibody specifically recognizes tenascin W having the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4, for use as a pharmaceutical.

30

14. The use of an antibody that specifically recognizes tenascin W for the manufacture of a medicament, for the prophylaxis or treatment of cancer.

15. The use as claimed in claim 14, wherein said cancer is metastatic.

16. The use as claimed in claim 14 or 15, wherein the cancer is a solid tumour.

17. The use as claimed in any of claims 14 to 16, wherein the cancer is a glioblastoma, prostate, lung, colorectal, osteo- or breast carcinoma.

5 18. The use of an antibody that specifically recognizes tenascin W for the prophylaxis or treatment of a bone disease resulting from excessive bone growth.

19. A composition comprising an isolated nucleic acid molecule having a nucleotide sequence selected from the group consisting of:

- 10 (a) a nucleotide sequence as set forth in SEQ ID No. 1 or SEQ ID No. 3;
- (b) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO: 4;
- (c) a nucleotide sequence with at least 35% identity to any
- 15 one of the sequences of (a) or (b);
- (d) a subsequence of a least 15 consecutive nucleotides of the sequence of (a), (b) or (c); and
- (e) a nucleotide sequence complementary to (a), (b), (c), or (d),

20 and a pharmaceutically acceptable excipient, diluent or carrier.

20. The composition of claim 19 wherein said nucleic acid molecule has a subsequence that is antisense to SEQ ID NO:1 or SEQ ID NO:3.

25 21. The composition of claim 19, said composition comprising said isolated nucleic acid molecule encoding the amino acid sequence shown in SEQ ID NO: 2 or SEQ ID NO:4.

30 22. A composition as claimed in any one of claims 19 to 21 for use as a pharmaceutical.

23. The use of the compositions of any of claims 19 to 21 for the manufacture of a medicament for the prophylaxis or treatment of cancer.

24. The use of the compositions of any of claims 19 to 21 for the manufacture of a medicament for the prophylaxis or treatment of bone pathologies.

5

25. A composition comprising tenascin-W and a pharmaceutically acceptable excipient, diluent or carrier.

26. The composition of claim 25, wherein said tenascin-W is recombinant.

10

27. A composition as claimed in claim 25, wherein said tenascin-W is a polypeptide having:

(a) an amino acid sequence as set forth in SEQ ID No. 2 or 4;

15

(b) an amino acid sequence with at least 35% identity to the sequence of (a); or  
a subsequence of at least 30 consecutive amino acids of the sequence of (a) or (b).

20

28. The composition of claim 27, said composition comprising said polypeptide encoding the amino acid sequence shown in SEQ ID NO: 4.

25

29. A composition as claimed in any one of claims 25 to 28 for use as a pharmaceutical.

30

30. The use of the compositions of any one of claims 25 to 28 for the manufacture of a medicament for the treatment or prophylactic treatment of any disease or condition requiring increased tenascin-W levels, e.g. thrombosis, wound healing or atherosclerosis, or for the treatment or prophylactic treatment of a condition ameliorated by the promotion of osteogenesis, e.g. bone healing, osteoporosis.



- 5 31. A method for treatment or prophylactic treatment of a disease or condition requiring increased tenascin-W levels, e.g. thrombosis, wound healing, atherosclerosis, bone healing and osteoporosis, said method comprising administering an effective amount of the compositions of any one of claims 25 to 28 to an individual in need of such treatment.
- 10 32. A method of inducing stem cell differentiation into bone cells, said method comprising contacting a suitable stem cell with an effective amount of the composition of any one of claims 25 to 28.
33. The method of claim 32, wherein said stem cell is a mesenchymal stem cell.
- 15 34. The use of tenascin-W as a stem cell marker.
35. A method for identifying modulators of tenascin W function, said method comprising contacting a test compound with a tenascin-W expressing cell and then measuring a change in one or more of:
- 20 (a) cell proliferation, e.g. cell progression entering S-phase of the cell cycle;
- (b) DNA synthesis;
- (c) cell adhesion;
- (d) cell spreading;
- 25 (e) focal adhesion and actin stress fibre formation on fibronectin; or
- (f) cell binding to extracellular matrix (ECM)
- relative to when said test compound is absent.
- 30 36. A method as claimed in claim 35, further comprising measuring a change in tenascin-W expression in the absence and presence of said test compound.

37. A method as claimed in claim 35 or 36, wherein said cell is a human cell.

5 38. A method as claimed in any of claims 35 to 37, wherein said cell is present in a tissue sample.

39. A method as claimed in any of claims 35 to 37, wherein said cell is present in a blood sample.

10 40. A method for identifying modulators of tenascin W function, said method comprising:

- (a) contacting a test compound with tenascin W and/or alpha8 beta1 integrin, and
- (b) measuring the binding of said test compound to tenascin-W and/or alpha8 beta1 integrin, or
- 15 (c) measuring a disruption of binding of tenascin-W to alpha8 beta1 integrin,

relative to when said test compound is absent.

20 41. A method as claimed in claim 40, further comprising measuring the binding of a control compound to tenascin-W.

42. A method as claimed in claim 40 or 41, wherein said tenascin-W is attached to a solid surface.

25 43. A method as claimed in any one of claims 41 to 42, wherein said binding is detected using an antibody labelled with a fluorescent label, a fluorescence quencher, a radioactive label, a scintillant or an enzyme.

30 44. A method as claimed in any one of claims 41 to 43, wherein a decrease in binding of tenascin-W to alpha8 beta1 integrin is indicative of an inhibitor of tenascin W function.

45. A substance for the prevention or the prophylactic treatment of a disease or condition dependent on tenascin-W, identified by a method as claimed in any of claims 40 to 44.

- 5      46. A method of diagnosing or prognosing cancer comprising:
- (a)      analysing a sample obtained from an individual for the presence of tenascin-W protein or transcript
  - (b)      correlating the presence of tenascin-W with an unfavourable prognosis or diagnosis.

10

47. A method of diagnosing or prognosing cancer as claimed in claim 46, comprising correlating in (b) an elevated level of tenascin-W protein or transcript relative to healthy tissue with an unfavourable prognosis or diagnosis.

15

48. A method of diagnosing or prognosing cancer as claimed in any of claims 46 to 47, wherein said sample is blood serum or plasma from an individual.

20

49. A method of diagnosing or prognosing cancer as claimed in any of claims 46 to 48, further comprising analysing said sample for the presence of alpha 8 integrin expression, the presence of said alpha 8 integrin correlating with an unfavourable prognosis or diagnosis.

25

50. A method of diagnosing or prognosing cancer as claimed in any of claims 46 to 49, further comprising propagating cells in said sample in cell culture.

30

51. A method of diagnosing or prognosing cancer as claimed in any of claims 48 to 50, wherein said tenascin-W protein is detected using an antibody specific for tenascin-W.

52. A method of diagnosing or prognosing cancer as claimed in any of claims 48 to 50, wherein said tenascin-W transcripts are detected using a polymerase chain reaction.

## SEQUENCE LISTING

**<110> Novartis Forschungsstiftung, Zweigniederlassung Friedrich Miescher  
Institute for Biomedical Research**

## **<120> Tenascin-W Compositions and Uses thereof**

<130> 1-32411/P1/FMI

<160> 16

<170> PatentIn version 3.1

**<210> 1**

**<211> 3891**

**<212> DNA**

<213> Mus musculus

**<220>**

**<221> CDS**

<222> (1)..(3891)

**<223>**

<400> 1

atg ggt ctc tgg ggg atg ctc gcc ttc ccc ctg gga ttt ctg ctt gct 48  
Met Gly Leu Trp Gly Met Leu Ala Phe Pro Leu Gly Phe Leu Leu Ala  
1 5 10 15

tct gtg ctc ctg gtg gct tcg gcc cca gcc act cca gag tct ccc ggc 96  
Ser Val Leu Leu Val Ala Ser Ala Pro Ala Thr Pro Glu Ser Pro Gly  
20 25 30

tgc agc aac aaa gag caa cag gtc act gtt agc cac acc tac aag att 144  
Cys Ser Asn Lys Glu Gln Gln Val Thr Val Ser His Thr Tyr Lys Ile  
35 40 45

gac gtg ccc aag tct gct ctg gtt caa gta gag acc gac cca cag tca 192  
Asp Val Pro Lys Ser Ala Leu Val Gln Val Glu Thr Asp Pro Gln Ser  
50 55 60

ctc agc gat gat ggg aca tca ctc ttg gct ccc ggg gag gat ggg gag 240  
Leu Ser Asp Asp Gly Thr Ser Leu Leu Ala Pro Gly Glu Asp Gly Glu  
65 70 75 80

gag cag aac att atc ttc agg cac aac atc cgt ctt cag aca ccg cag 288  
Glu Gln Asn Ile Ile Phe Arg His Asn Ile Arg Leu Gln Thr Pro Gln

85	90	95	
aag aat tgc gac ctg gca gac agt gtc cag gac ctg cta gcc cgg atg Lys Asn Cys Asp Leu Ala Asp Ser Val Gln Asp Leu Leu Ala Arg Met 100 105 110			336
aaa aag ctg gag gaa gag atg gca gag ctg aag gag cag tgc aat acc Lys Lys Leu Glu Glu Glu Met Ala Glu Leu Lys Glu Gln Cys Asn Thr 115 120 125			384
aac cgc tgc tgc cag ggg gct gct gat ctg agc cgt cac tgc agt ggc Asn Arg Cys Cys Gln Gly Ala Ala Asp Leu Ser Arg His Cys Ser Gly 130 135 140			432
cac ggg acc ttc ctc cct gag acc tgc agc tgc cac tgt gac cag ggc His Gly Thr Phe Leu Pro Glu Thr Cys Ser Cys His Cys Asp Gln Gly 145 150 155 160			480
tgg gag ggc gca gac tgt gat cag ccc acc tgt cct ggg gct tgc aac Trp Glu Gly Ala Asp Cys Asp Gln Pro Thr Cys Pro Gly Ala Cys Asn 165 170 175			528
ggc cac ggg cgc tgt gtg gat ggg cag tgc gtg tgt gac ggc ccc cac Gly His Gly Arg Cys Val Asp Gly Gln Cys Val Cys Asp Ala Val Thr 180 185 190			576
gtg ggg gtc gac tgc gcc tac gcc gcc tgt ccc cag gac tgc agt ggc Val Gly Val Asp Cys Ala Tyr Ala Ala Cys Pro Gln Asp Cys Ser Gly 195 200 205			624
cat ggc gtg tgc gtg cag ggt gtc tgc cag tgc cac gag gac ttc aca His Gly Val Cys Val Gln Gly Val Cys Gln Cys His Glu Asp Phe Thr 210 215 220			672
gca gag gac tgc agc gag cag cgc tgt cct ggc gac tgt agt ggc aat Ala Glu Asp Cys Ser Glu Gln Arg Cys Pro Gly Asp Cys Ser Gly Asn 225 230 235 240			720
ggt ttc tgt gac act ggc gag tgt tac tgt gag atg ggc ttt act ggc Gly Phe Cys Asp Thr Gly Glu Cys Tyr Cys Glu Met Gly Phe Thr Gly 245 250 255			768
ccc gac tgt tcc cag gtg gtg gct cct cag ggc ctg cag ttg ctc aag Pro Asp Cys Ser Gln Val Val Ala Pro Gln Gly Leu Gln Leu Leu Lys 260 265 270			816

agc acg gag aac tct ctg ctg gtg agt tgg gag ccc tcc agt gag gta Ser Thr Glu Asn Ser Leu Leu Val Ser Trp Glu Pro Ser Ser Glu Val 275 280 285	864
gac tac tac ctg ctc agc tac tac ccc ctg ggg aag gag caa gct aca Asp Tyr Tyr Leu Leu Ser Tyr Tyr Pro Leu Gly Lys Glu Gln Ala Thr 290 295 300	912
aaa cag gtc cgg gta ccc aag gag cag cac acc tat gac atc acc ggc Lys Gln Val Arg Val Pro Lys Glu Gln His Thr Tyr Asp Ile Thr Gly 305 310 315 320	960
ttg ctg cct gga acc aag tac ata gtc acc ctg cgc aac gtg aag aaa Leu Leu Pro Gly Thr Lys Tyr Ile Val Thr Leu Arg Asn Val Lys Lys 325 330 335	1008
gac att tcc agc agc cct cag cat cta ctt gcc acc aca gac ctt gct Asp Ile Ser Ser Ser Pro Gln His Leu Leu Ala Thr Thr Asp Leu Ala 340 345 350	1056
gtg ctt ggc act gcc tgg gta aat gaa gac act gag aca tcc ctc gat Val Leu Gly Thr Ala Trp Val Asp Glu Thr Glu Thr Ser Leu Asp 355 360 365	1104
gtg gag tgg gag aac cct ctg act gag gtc gac tat tac aag ctt cgg Val Glu Trp Glu Asn Pro Leu Thr Glu Val Asp Tyr Tyr Lys Leu Arg 370 375 380	1152
tat ggc ccc tta aca ggg cag gag gtg aca gag gtc act gtg ccc aag Tyr Gly Pro Leu Thr Gly Gln Glu Val Thr Glu Val Thr Val Pro Lys 385 390 395 400	1200
agc cgt gat ccc aag agc aga tat gac atc act ggt ctg cag cct gga Ser Arg Asp Pro Lys Ser Arg Tyr Asp Ile Thr Gly Leu Gln Pro Gly 405 410 415	1248
acg gaa tat aaa atc aca gtt gtg ccc atc cga ggt gat ctg gag gga Thr Glu Tyr Lys Ile Thr Val Val Pro Ile Arg Gly Asp Leu Glu Gly 420 425 430	1296
aag ccg att ctc ctg aat ggc agg aca gaa att gat gga cca acc aat Lys Pro Ile Leu Leu Asn Gly Arg Thr Glu Ile Asp Gly Pro Thr Asn 435 440 445	1344
gtg gtc aca aat cag gtg aca gaa gac aca gca tct gtt tcc tgg gat Val Val Thr Asn Gln Val Thr Glu Asp Thr Ala Ser Val Ser Trp Asp 450 455 460	1392

450	455	460	
cca gtg agg gct gac ata gac aag tat gtg gtg cgc tat atc gcc ccc			1440
Pro Val Arg Ala Asp Ile Asp Lys Tyr Val Val Arg Tyr Ile Ala Pro			
465	470	475	480
gat ggg gag acc aag gag aag gca gta cca aag gac cag agc agc acc			1488
Asp Gly Glu Thr Lys Glu Lys Ala Val Pro Lys Asp Gln Ser Ser Thr			
	485	490	495
gtt ctc aca ggc ctg aag cca gga gag gcc tac aaa gtc ttt gtg tgg			1536
Val Leu Thr Gly Leu Lys Pro Gly Glu Ala Tyr Lys Val Phe Val Trp			
	500	505	510
gct gag agg ggc aac caa ggc agc aag aaa gca gac acc aag gcc ctc			1584
Ala Glu Arg Gly Asn Gln Gly Ser Lys Lys Ala Asp Thr Lys Ala Leu			
	515	520	525
aca gaa att gac agt cca gaa aac ctg gtg act gac cgg gtg aca gag			1632
Thr Glu Ile Asp Ser Pro Glu Asn Leu Val Thr Asp Arg Val Thr Glu			
	530	535	540
aac agc gca tat gtc tgg ggg gac cca gtg gag gct gac atc gac agg			1680
Asn Ser Leu Ser Val Trp Asp Pro Val Glu Ala Asp Ile Asp Arg			
545	550	555	560
tat gtg gta agc tac act tcc gtg gat gga gag acg aag cag gtt cca			1728
Tyr Val Val Ser Tyr Thr Ser Val Asp Gly Glu Thr Lys Gln Val Pro			
	565	570	575
gtg aag aag gac cag agg agc acc gtc ctc acc ggc ctg agt ccc ggt			1776
Val Lys Lys Asp Gln Arg Ser Thr Val Leu Thr Gly Leu Ser Pro Gly			
	580	585	590
gtg gag tac aaa gtt tac gtg tgg gca gag aaa ggc gat cgg gag agc			1824
Val Glu Tyr Lys Val Tyr Val Trp Ala Glu Lys Gly Asp Arg Glu Ser			
	595	600	605
aag aag gcc aac acc aag gct ccc aca gac atc gac agc ccc aaa aac			1872
Lys Lys Ala Asn Thr Lys Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn			
	610	615	620
ttg gta act gac cag gtg aca gag aac act ctc agt gtc tcc tgg gac			1920
Leu Val Thr Asp Gln Val Thr Glu Asn Thr Leu Ser Val Ser Trp Asp			
625	630	635	640



cct gtt cag gcc aac att gac agg tat atg gtg agc tac acc tct gcc Pro Val Gln Ala Asn Ile Asp Arg Tyr Met Val Ser Tyr Thr Ser Ala 645 650 655	1968
gat gga gag aca aga gag gtc cca gtg cct aag gag aag agc agt acc Asp Gly Glu Thr Arg Glu Val Pro Val Pro Lys Glu Lys Ser Ser Thr 660 665 670	2016
gtc ctg act ggc ctg agg cca ggt gtg gag tac aag gtc cat gtg tgg Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Lys Val His Val Trp 675 680 685	2064
gcc cag aag ggg acc cag gag agc aga aag gcc aac acc aag gcc ccc Ala Gln Lys Gly Thr Gln Glu Ser Arg Lys Ala Asn Thr Lys Ala Pro 690 695 700	2112
aca gat att gat ggc ccc aaa aac ctg gtg act gac cag gtg acc gag Thr Asp Ile Asp Gly Pro Lys Asn Leu Val Thr Asp Gln Val Thr Glu 705 710 715 720	2160
acc act ctt agt gtc tcc tgg gac cca gtg gag gct gac att gat agg Thr Thr Leu Ser Val Ser Trp Asp Pro Val Glu Ala Asp Ile Asp Arg 725 730 735	2208
tac atg gtt cgc tac acg tct cct gat gga gag acc aag gag gtg cct Tyr Met Val Arg Tyr Thr Ser Pro Asp Gly Glu Thr Lys Glu Val Pro 740 745 750	2256
gtg tca aag gat aag agc agc aca gtc ctg agg ggc ctg agg cca ggt Val Ser Lys Asp Lys Ser Ser Thr Val Leu Arg Gly Leu Arg Pro Gly 755 760 765	2304
gtg gag tac aag gtg gat gta tgg gcc cag aag ggg gcc cag gac agc Val Glu Tyr Lys Val Asp Val Trp Ala Gln Lys Gly Ala Gln Asp Ser 770 775 780	2352
cgg aag gcc aac acc aag gcc ccc aca gat att gac agc cct aaa aac Arg Lys Ala Asn Thr Lys Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn 785 790 795 800	2400
cta gtg act gag cag gtg gca gag agc act gcc acc gtg tcc tgg gac Leu Val Thr Glu Gln Val Ala Glu Ser Thr Ala Thr Val Ser Trp Asp 805 810 815	2448
cca gta gag gct gac atc gac agg tat gtg gtg cgc tac acc tct gct Pro Val Glu Ala Asp Ile Asp Arg Tyr Val Val Arg Tyr Thr Ser Ala 820 825 830 835	2496

820	825	830	
gat gga gag acc agg gag att cca gtg agg aag gag aag agc agc act			2544
Asp Gly Glu Thr Arg Glu Ile Pro Val Arg Lys Glu Lys Ser Ser Thr			
835	840	845	
gtc ctc aca ggc ctg aga ccg ggt gtg gag tac acg gtc caa gtg tgg			2592
Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Thr Val Gln Val Trp			
850	855	860	
gct cag aag ggg gcc cgg gag agc aag aag gcc aaa acc aag gcc ccc			2640
Ala Gln Lys Gly Ala Arg Glu Ser Lys Lys Ala Lys Thr Lys Ala Pro			
865	870	875	880
aca gaa att gac agc ccc aag aac ttg gtg acc aac cga gtg aca gaa			2688
Thr Glu Ile Asp Ser Pro Lys Asn Leu Val Thr Asn Arg Val Thr Glu			
885	890	895	
aat aca gcc acc atc tcc tgg gac cca gtg cga gcc aac att gac agg			2736
Asn Thr Ala Thr Ile Ser Trp Asp Pro Val Arg Ala Asn Ile Asp Arg			
900	905	910	
tac atg gtt cgc tac acc tct gcg gat gga gag act aag gag aca cca			2784
Tyr Met Val Arg Tyr Thr Ser Ala Asp Gly Glu Thr Lys Glu Ala Pro			
915	920	925	
gtg tca aag gat cag agt aac acc atc ctg aca ggc ctg aaa cca ggc			2832
Val Ser Lys Asp Gln Ser Asn Thr Ile Leu Thr Gly Leu Lys Pro Gly			
930	935	940	
atg gaa tat acc att cat gtg tgg gcc cag aag ggg gcc cgg gag agc			2880
Met Glu Tyr Thr Ile His Val Trp Ala Gln Lys Gly Ala Arg Glu Ser			
945	950	955	960
aag aag gct gat acc aag gcc cta aca gaa att gac cct ccc aga aat			2928
Lys Lys Ala Asp Thr Lys Ala Leu Thr Glu Ile Asp Pro Pro Arg Asn			
965	970	975	
ctc cgt ccg ttc ggg gta aca cat tct ggt ggg gtt ttg acc tgg ttg			2976
Leu Arg Pro Phe Gly Val Thr His Ser Gly Gly Val Leu Thr Trp Leu			
980	985	990	
ccc cca tct gct caa att gat ggc tac att ttg acc tac cag ttc cca			3024
Pro Pro Ser Ala Gln Ile Asp Gly Tyr Ile Leu Thr Tyr Gln Phe Pro			
995	1000	1005	

aat ggc acc gtg aag ggg gtg gag ctc cca aga ggc cag cag aga	3069
Asn Gly Thr Val Lys Gly Val Glu Leu Pro Arg Gly Gln Gln Arg	
1010 1015 1020	
ttt gaa ttg caa gac ctg gaa cag ggt gtc acc tat cct gtt tcc	3114
Phe Glu Leu Gln Asp Leu Glu Gln Gly Val Thr Tyr Pro Val Ser	
1025 1030 1035	
ttg gtt gcc ttc aaa ggt aat cag cgg agc cgg act gtg tct acc	3159
Leu Val Ala Phe Lys Gly Asn Gln Arg Ser Arg Thr Val Ser Thr	
1040 1045 1050	
acc ctt tct aca gtg gat gct cgc ttt cca cac ccc tca gac tgc	3204
Thr Leu Ser Thr Val Asp Ala Arg Phe Pro His Pro Ser Asp Cys	
1055 1060 1065	
agt caa gtt cag cag aac acc aac gct gcc agt ggg ctc tac acc	3249
Ser Gln Val Gln Gln Asn Thr Asn Ala Ala Ser Gly Leu Tyr Thr	
1070 1075 1080	
atc tac ctc aat ggt gat gcc agc cgg ctc cag gtg tac tgc	3294
Ile Tyr Leu Asn Gly Asp Ala Ser Arg Pro Met Glu Val Tyr Cys	
1085 1090 1095	
gac atg gac act gac gga ggc ggc tgg gtc gtc ttc cag aga cgg	3339
Asp Met Asp Thr Asp Gly Gly Gly Trp Ile Val Phe Gln Arg Arg	
1100 1105 1110	
aac act ggg cag ctg gat ttc ttc aag cgt tgg cgg agt tat gta	3384
Asn Thr Gly Gln Leu Asp Phe Phe Lys Arg Trp Arg Ser Tyr Val	
1115 1120 1125	
gaa ggt ttt ggg gac ccc atg aag gag ttc tgg ctt gga ctt gat	3429
Glu Gly Phe Gly Asp Pro Met Lys Glu Phe Trp Leu Gly Leu Asp	
1130 1135 1140	
aaa cta cat aat ctc acc act ggc acc acc act cgg tat gag gtg	3474
Lys Leu His Asn Leu Thr Thr Gly Thr Thr Thr Arg Tyr Glu Val	
1145 1150 1155	
agg gca gac tta cag act ttc aat gaa tct gcc tac gct gta tat	3519
Arg Ala Asp Leu Gln Thr Phe Asn Glu Ser Ala Tyr Ala Val Tyr	
1160 1165 1170	
gat ttc ttc caa gtg gca tcc agc aaa gag cgg tac aag ctg tcg	3564
Asp Phe Phe Gln Val Ala Ser Ser Lys Glu Arg Tyr Lys Leu Ser	

1175	1180	1185	
gtt ggg aaa tac aga ggc aca	gcc ggg gat gct ctc	acc tac cac	3609
Val Gly Lys Tyr Arg Gly Thr	Ala Gly Asp Ala Leu	Thr Tyr His	
1190	1195	1200	
aat gga tgg aag ttc acg act	ttt gac aga gac agt	gat att gcc	3654
Asn Gly Trp Lys Phe Thr Thr	Phe Asp Arg Asp Ser	Asp Ile Ala	
1205	1210	1215	
ctc agc aac tgt gca ctg acg	cat cat ggt ggc tgg	tgg tat aag	3699
Leu Ser Asn Cys Ala Leu Thr	His His Gly Gly Trp	Trp Tyr Lys	
1220	1225	1230	
aac tgc cat ttg gcc aac ccg	aat ggc aaa tat ggg	gag acc aag	3744
Asn Cys His Leu Ala Asn Pro	Asn Gly Lys Tyr Gly	Glu Thr Lys	
1235	1240	1245	
cac agc gag ggg gtg aac tgg	gag cca tgg aag gga	cat gag ttc	3789
His Ser Glu Gly Val Asn Trp	Glu Pro Trp Lys Gly	His Glu Phe	
1250	1255	1260	
tcc att cca tac gtg gag cgc	aaa atc cgc ccg ttt	ggg tac agc	3834
Ser Ile Pro Tyr Asn Leu	Lys Ile Arg Pro Phe	Gly Tyr Ser	
1265	1270	1275	
aga gac cgt ttc tct ggc aga	aag aag cgc tcc ata	gga aaa gca	3879
Arg Asp Arg Phe Ser Gly Arg	Lys Lys Arg Ser Ile	Gly Lys Ala	
1280	1285	1290	
agg atg ttc tga			3891
Arg Met Phe			
1295			

&lt;210&gt; 2

&lt;211&gt; 1296

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 2

Met Gly Leu Trp Gly Met Leu Ala Phe Pro Leu Gly Phe Leu Leu Ala  
 1 5 10 15

Ser Val Leu Leu Val Ala Ser Ala Pro Ala Thr Pro Glu Ser Pro Gly

20	25	30
Cys Ser Asn Lys Glu Gln Gln Val Thr Val Ser His Thr Tyr Lys Ile		
35	40	45
Asp Val Pro Lys Ser Ala Leu Val Gln Val Glu Thr Asp Pro Gln Ser		
50	55	60
Leu Ser Asp Asp Gly Thr Ser Leu Leu Ala Pro Gly Glu Asp Gly Glu		
65	70	75
Glu Gln Asn Ile Ile Phe Arg His Asn Ile Arg Leu Gln Thr Pro Gln		
85	90	95
Lys Asn Cys Asp Leu Ala Asp Ser Val Gln Asp Leu Leu Ala Arg Met		
100	105	110
Lys Lys Leu Glu Glu Glu Met Ala Glu Leu Lys Glu Gln Cys Asn Thr		
115	120	125
En Arg Cys Cys Gln Gly Ala Ala Asp Leu Ser Arg His Cys Ser Gly		
130	135	140
Gly Thr Phe Leu Pro Glu Thr Cys Ser Cys His Cys Asp Gln Gly		
145	150	155
Trp Glu Gly Ala Asp Cys Asp Gln Pro Thr Cys Pro Gly Ala Cys Asn		
165	170	175
Gly His Gly Arg Cys Val Asp Gly Gln Cys Val Cys Asp Ala Pro Tyr		
180	185	190
Val Gly Val Asp Cys Ala Tyr Ala Ala Cys Pro Gln Asp Cys Ser Gly		
195	200	205
His Gly Val Cys Val Gln Gly Val Cys Gln Cys His Glu Asp Phe Thr		
210	215	220
Ala Glu Asp Cys Ser Glu Gln Arg Cys Pro Gly Asp Cys Ser Gly Asn		
225	230	235
Gly Phe Cys Asp Thr Gly Glu Cys Tyr Cys Glu Met Gly Phe Thr Gly		
245	250	255
Pro Asp Cys Ser Gln Val Val Ala Pro Gln Gly Leu Gln Leu Leu Lys		
260	265	270

Ser Thr Glu Asn Ser Leu Leu Val Ser Trp Glu Pro Ser Ser Glu Val  
275 280 285

Asp Tyr Tyr Leu Leu Ser Tyr Tyr Pro Leu Gly Lys Glu Gln Ala Thr  
290 295 300

Lys Gln Val Arg Val Pro Lys Glu Gln His Thr Tyr Asp Ile Thr Gly  
305 310 315 320

Leu Leu Pro Gly Thr Lys Tyr Ile Val Thr Leu Arg Asn Val Lys Lys  
325 330 335

Asp Ile Ser Ser Ser Pro Gln His Leu Leu Ala Thr Thr Asp Leu Ala  
340 345 350

Val Leu Gly Thr Ala Trp Val Asn Glu Glu Thr Glu Thr Ser Leu Asp  
355 360 365

Val Glu Trp Glu Asn Pro Leu Thr Glu Val Asp Tyr Tyr Lys Leu Arg  
370 375 380

Tyr Gly Pro Leu Thr Gly Gln Glu Val Thr Glu Val Thr Val Pro  
385 390 395

Ser Arg Asp Pro Lys Ser Arg Tyr Asp Ile Thr Gly Leu Gln Pro Gly  
405 410 415

Thr Glu Tyr Lys Ile Thr Val Val Pro Ile Arg Gly Asp Leu Glu Gly  
420 425 430

Lys Pro Ile Leu Leu Asn Gly Arg Thr Glu Ile Asp Gly Pro Thr Asn  
435 440 445

Val Val Thr Asn Gln Val Thr Glu Asp Thr Ala Ser Val Ser Trp Asp  
450 455 460

Pro Val Arg Ala Asp Ile Asp Lys Tyr Val Val Arg Tyr Ile Ala Pro  
465 470 475 480

Asp Gly Glu Thr Lys Glu Lys Ala Val Pro Lys Asp Gln Ser Ser Thr  
485 490 495

Val Leu Thr Gly Leu Lys Pro Gly Glu Ala Tyr Lys Val Phe Val Trp  
500 505 510

Ala Glu Arg Gly Asn Gln Gly Ser Lys Lys Ala Asp Thr Lys Ala Leu  
515 520 525

Thr Glu Ile Asp Ser Pro Glu Asn Leu Val Thr Asp Arg Val Thr Glu  
530 535 540

Asn Ser Leu Ser Val Ser Trp Asp Pro Val Glu Ala Asp Ile Asp Arg  
545 550 555 560

Tyr Val Val Ser Tyr Thr Ser Val Asp Gly Glu Thr Lys Gln Val Pro  
565 570 575

Val Lys Lys Asp Gln Arg Ser Thr Val Leu Thr Gly Leu Ser Pro Gly  
580 585 590

Val Glu Tyr Lys Val Tyr Val Trp Ala Glu Lys Gly Asp Arg Glu Ser  
595 600 605

Lys Lys Ala Asn Thr Lys Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn  
610 615 620

Leu Val Thr Asp Gln Val Thr Glu Asn Thr Leu Ser Val Ser Trp Asp  
625 630 635 640

Pro Val Gln Ala Asn Ile Asp Arg Tyr Met Val Ser Tyr Thr Ser Ala  
645 650 655

Asp Gly Glu Thr Arg Glu Val Pro Val Pro Lys Glu Lys Ser Ser Thr  
660 665 670

Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Lys Val His Val Trp  
675 680 685

Ala Gln Lys Gly Thr Gln Glu Ser Arg Lys Ala Asn Thr Lys Ala Pro  
690 695 700

Thr Asp Ile Asp Gly Pro Lys Asn Leu Val Thr Asp Gln Val Thr Glu  
705 710 715 720

Thr Thr Leu Ser Val Ser Trp Asp Pro Val Glu Ala Asp Ile Asp Arg  
725 730 735

Tyr Met Val Arg Tyr Thr Ser Pro Asp Gly Glu Thr Lys Glu Val Pro  
740 745 750

Val Ser Lys Asp Lys Ser Ser Thr Val Leu Arg Gly Leu Arg Pro Gly  
755 760 765

Val Glu Tyr Lys Val Asp Val Trp Ala Gln Lys Gly Ala Gln Asp Ser  
770 775 780

Arg Lys Ala Asn Thr Lys Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn  
785 790 795 800

Leu Val Thr Glu Gln Val Ala Glu Ser Thr Ala Thr Val Ser Trp Asp  
805 810 815

Pro Val Glu Ala Asp Ile Asp Arg Tyr Val Val Arg Tyr Thr Ser Ala  
820 825 830

Asp Gly Glu Thr Arg Glu Ile Pro Val Arg Lys Glu Lys Ser Ser Thr  
835 840 845

Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Thr Val Gln Val Trp  
850 855 860

Ala Gln Lys Gly Ala Arg Glu Ser Lys Lys Ala Lys Thr Lys Ala Pro  
865 870 875 880

Thr Glu Ile Asp Ser Pro Lys Asn Leu Val Thr Asn Arg Val Thr Glu  
885 890 895

Asn Thr Ala Thr Ile Ser Trp Asp Pro Val Arg Ala Asn Ile Asp Arg  
900 905 910

Tyr Met Val Arg Tyr Thr Ser Ala Asp Gly Glu Thr Lys Glu Ile Pro  
915 920 925

Val Ser Lys Asp Gln Ser Asn Thr Ile Leu Thr Gly Leu Lys Pro Gly  
930 935 940

Met Glu Tyr Thr Ile His Val Trp Ala Gln Lys Gly Ala Arg Glu Ser  
945 950 955 960

Lys Lys Ala Asp Thr Lys Ala Leu Thr Glu Ile Asp Pro Pro Arg Asn  
965 970 975

Leu Arg Pro Phe Gly Val Thr His Ser Gly Gly Val Leu Thr Trp Leu  
980 985 990



Pro Pro Ser Ala Gln Ile Asp Gly Tyr Ile Leu Thr Tyr Gln Phe Pro  
 995 1000 1005

Asn Gly Thr Val Lys Gly Val Glu Leu Pro Arg Gly Gln Gln Arg  
 1010 1015 1020

Phe Glu Leu Gln Asp Leu Glu Gln Gly Val Thr Tyr Pro Val Ser  
 1025 1030 1035

Leu Val Ala Phe Lys Gly Asn Gln Arg Ser Arg Thr Val Ser Thr  
 1040 1045 1050

Thr Leu Ser Thr Val Asp Ala Arg Phe Pro His Pro Ser Asp Cys  
 1055 1060 1065

Ser Gln Val Gln Gln Asn Thr Asn Ala Ala Ser Gly Leu Tyr Thr  
 1070 1075 1080

Ile Tyr Leu Asn Gly Asp Ala Ser Arg Pro Met Gln Val Tyr Cys  
 1085 1090 1095

Asp Met Asp Thr Asp Gly Gly Gly Trp Ile Val Phe Gln Arg Arg  
 1100 1105 1110

Asp Thr Gly Gln Leu Asp Phe Phe Lys Arg Trp Arg Ser Tyr Val  
 1115 1120 1125

Glu Gly Phe Gly Asp Pro Met Lys Glu Phe Trp Leu Gly Leu Asp  
 1130 1135 1140

Lys Leu His Asn Leu Thr Thr Gly Thr Thr Thr Arg Tyr Glu Val  
 1145 1150 1155

Arg Ala Asp Leu Gln Thr Phe Asn Glu Ser Ala Tyr Ala Val Tyr  
 1160 1165 1170

Asp Phe Phe Gln Val Ala Ser Ser Lys Glu Arg Tyr Lys Leu Ser  
 1175 1180 1185

Val Gly Lys Tyr Arg Gly Thr Ala Gly Asp Ala Leu Thr Tyr His  
 1190 1195 1200

Asn Gly Trp Lys Phe Thr Thr Phe Asp Arg Asp Ser Asp Ile Ala  
 1205 1210 1215

Leu Ser Asn Cys Ala Leu Thr His His Gly Gly Trp Trp Tyr Lys

1220	1225	1230
Asn Cys His Leu Ala Asn Pro Asn Gly Lys Tyr Gly Glu Thr Lys		
1235	1240	1245
His Ser Glu Gly Val Asn Trp Glu Pro Trp Lys Gly His Glu Phe		
1250	1255	1260
Ser Ile Pro Tyr Val Glu Leu Lys Ile Arg Pro Phe Gly Tyr Ser		
1265	1270	1275
Arg Asp Arg Phe Ser Gly Arg Lys Lys Arg Ser Ile Gly Lys Ala		
1280	1285	1290
Arg Met Phe		
1295		

<210> 3  
 <211> 3885  
 <212> DNA  
 <213> homo sapiens

<220>  
 <221> CDS  
 <222> (1)..(3885)  
 <223>

<300>  
 <308> AL049689  
 <309> 1999-04-21  
 <313> (1)..(3885)

<400> 3

atg ttc cgc ttc cct atg ggg ctc ctg ctt ggc tct gtg ctc ctg gtg	48
Met Phe Arg Phe Pro Met Gly Leu Leu Leu Gly Ser Val Leu Leu Val	
1 5 10 15	
gct tcg gcc cca gcc act ctg gag cct ccc ggc tgc agc aac aag gag	96
Ala Ser Ala Pro Ala Thr Leu Glu Pro Pro Gly Cys Ser Asn Lys Glu	
20 25 30	
caa cag gtc act gtc agc cac acc tac aag atc gat gtg ccc aag tct	144
Gln Gln Val Thr Val Ser His Thr Tyr Lys Ile Asp Val Pro Lys Ser	
35 40 45	

gcc ttg gtt cag gtt gac gct gac cct cag ccc ctc agt gac gat ggg 192  
 Ala Leu Val Gln Val Asp Ala Asp Pro Gln Pro Leu Ser Asp Asp Gly  
 50 55 60

gct tcg ctc ttg gcc ctg ggg gag gcc agg gag gaa cag aac atc atc 240  
 Ala Ser Leu Leu Ala Leu Gly Glu Ala Arg Glu Glu Gln Asn Ile Ile  
 65 70 75 80

ttc agg cac aac atc cgc ctt cag acg cca cag aag gac tgc gag ttg 288  
 Phe Arg His Asn Ile Arg Leu Gln Thr Pro Gln Lys Asp Cys Glu Leu  
 85 90 95

gca ggc agt gtc cag gac ctc ctg gcc cgg gtg aag aag ctg gag gaa 336  
 Ala Gly Ser Val Gln Asp Leu Leu Ala Arg Val Lys Lys Leu Glu Glu  
 100 105 110

gag atg gtg gag atg aag gaa cag tgt agt gcc cag cgc tgc tgc cag 384  
 Glu Met Val Glu Met Lys Glu Gln Cys Ser Ala Gln Arg Cys Cys Gln  
 115 120 125

gga gtc act gat cta agc cgc cac tgc agc ggc cag cgc tgc ttc tcc 432  
 Gly Val Thr Asp Leu Ser Arg His Cys Ser Gly His Gly Thr Phe Ser  
 130 135 140

ctg gag acc tgc agc tgc cac tgc gaa gag ggc agg ggc ccc gcc 480  
 Leu Glu Thr Cys Ser Cys His Cys Glu Glu Gly Arg Glu Gly Pro Ala  
 145 150 155 160

tgc gag cgg ctg gcc tgc ccc ggg gcg tgc agc ggc cac ggg cgt tgc 528  
 Cys Glu Arg Leu Ala Cys Pro Gly Ala Cys Ser Gly His Gly Arg Cys  
 165 170 175

gtg gac ggg cgc tgc ctg tgc cat gag ccc tac gtg ggt gcc gac tgc 576  
 Val Asp Gly Arg Cys Leu Cys His Glu Pro Tyr Val Gly Ala Asp Cys  
 180 185 190

ggc tac ccg gcc tgc cct gag aac tgc agc gga cac ggc gag tgc gtg 624  
 Gly Tyr Pro Ala Cys Pro Glu Asn Cys Ser Gly His Gly Glu Cys Val  
 195 200 205

cgc ggc gtg tgc cag tgc cac gaa gac ttc atg tcg gag gac tgc agc 672  
 Arg Gly Val Cys Gln Cys His Glu Asp Phe Met Ser Glu Asp Cys Ser  
 210 215 220

gag aag cgc tgt ccc ggc gac tgc agc ggc cac ggc ttc tgt gac acg 720  
 Glu Lys Arg Cys Pro Gly Asp Cys Ser Gly His Gly Phe Cys Asp Thr

225	230	235	240	
ggc gag tgc tac tgc gag gag ggc ttc aca ggc ctg gac tgt gcc cag				768
Gly Glu Cys Tyr Cys Glu Glu Gly Phe Thr Gly Leu Asp Cys Ala Gln				
	245	250	255	
gtg gtc acc cca cag ggc ctg cag ctg ctc aag aac acg gag gat tct				816
Val Val Thr Pro Gln Gly Leu Gln Leu Leu Lys Asn Thr Glu Asp Ser				
	260	265	270	
ctg ctg gtg agc tgg gag ccc tcc agc cag gtg gat cac tac ctc ctc				864
Leu Leu Val Ser Trp Glu Pro Ser Ser Gln Val Asp His Tyr Leu Leu				
	275	280	285	
agc tac tac ccc ctg ggg aag gag ctc tct ggg aag cag atc caa gtg				912
Ser Tyr Tyr Pro Leu Gly Lys Glu Leu Ser Gly Lys Gln Ile Gln Val				
	290	295	300	
ccc aag gag cag cac agc tat gag att ctt ggt ttg ctg cct gga acc				960
Pro Lys Glu Gln His Ser Tyr Glu Ile Leu Gly Leu Leu Pro Gly Thr				
305	310	315	320	
aag tac ata gtc acc ctg cgc aac ctg aag aat gaa gtt tct agc agc				1008
Lys Tyr Ile Val Thr Leu Tyr Asn Lys Asn Glu Val Ser Ser Ser				
	325	330	335	
cca cag cat cta ctt gcc acc aca gac ctt gct gtg ctt ggc act gcc				1056
Pro Gln His Leu Leu Ala Thr Thr Asp Leu Ala Val Leu Gly Thr Ala				
	340	345	350	
tgg gtg aca gat gag act gag aac tcc ctt gac gtg gag tgg gaa aac				1104
Trp Val Thr Asp Glu Thr Glu Asn Ser Leu Asp Val Glu Trp Glu Asn				
	355	360	365	
ccc tca act gag gtg gac tac tac aag ctg cga tat ggc ccc atg aca				1152
Pro Ser Thr Glu Val Asp Tyr Tyr Lys Leu Arg Tyr Gly Pro Met Thr				
	370	375	380	
gga cag gag gta gct gag gtc act gtg ccc aag agc agt gac ccc aag				1200
Gly Gln Glu Val Ala Glu Val Thr Val Pro Lys Ser Ser Asp Pro Lys				
385	390	395	400	
agc cga tat gac atc act ggt ctg cac ccg ggg act gag tat aag atc				1248
Ser Arg Tyr Asp Ile Thr Gly Leu His Pro Gly Thr Glu Tyr Lys Ile				
	405	410	415	

acg gtg gtg ccc atg aga gga gag ctg gag ggc aag ccg atc ctc ctg	1296
Thr Val Val Pro Met Arg Gly Glu Leu Glu Gly Lys Pro Ile Leu Leu	
420 425 430	
aat ggc agg aca gaa att gac agt cca acc aat gtt gtc act gat cga	1344
Asn Gly Arg Thr Glu Ile Asp Ser Pro Thr Asn Val Val Thr Asp Arg	
435 440 445	
gtg act gaa gac aca gca act gtc tcc tgg gac cca gtg cag gct gtc	1392
Val Thr Glu Asp Thr Ala Thr Val Ser Trp Asp Pro Val Gln Ala Val	
450 455 460	
ata gac aag tat gta gtg cgc tac act tct gct gat ggg gac acc aag	1440
Ile Asp Lys Tyr Val Val Arg Tyr Thr Ser Ala Asp Gly Asp Thr Lys	
465 470 475 480	
gaa atg gca gtg cac aag gat gag agc agc act gtc ctg acg ggc ctg	1488
Glu Met Ala Val His Lys Asp Glu Ser Ser Thr Val Leu Thr Gly Leu	
485 490 495	
aag gca gaa gag gca tac aag gtc tac gtg tgg gct gaa agg ggc aac	1536
Asn Pro Gly Glu Ala Tyr Lys Val Tyr Val Trp Ala Glu Arg Gly Asn	
500 505 510	
cag agc aag aaa gct gac acc aat gcc ctc aca gaa att gac agc	1584
Gln Gly Ser Lys Lys Ala Asp Thr Asn Ala Leu Thr Glu Ile Asp Ser	
515 520 525	
cca gca aac ctg gtg act gac cgg gtg act gag aat acc gcc acc atc	1632
Pro Ala Asn Leu Val Thr Asp Arg Val Thr Glu Asn Thr Ala Thr Ile	
530 535 540	
tcc tgg gac ccg gta cag gcc acc att gac aag tac gtg gtg cgc tac	1680
Ser Trp Asp Pro Val Gln Ala Thr Ile Asp Lys Tyr Val Val Arg Tyr	
545 550 555 560	
acc tct gct gac gac caa gag acc aga gag gtt ctg gtg ggg aag gag	1728
Thr Ser Ala Asp Asp Gln Glu Thr Arg Glu Val Leu Val Gly Lys Glu	
565 570 575	
cag agc agc act gtc ctg aca ggc ctg agg cca ggt gtg gag tac aca	1776
Gln Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Thr	
580 585 590	
gtg cat gtc tgg gcc cag aag ggg gac cga gag agc aag aag gct gac	1824
Val His Val Trp Ala Gln Lys Gly Asp Arg Glu Ser Lys Lys Ala Asp	

595	600	605	
acc aac gcc ccg aca gat att gac agc ccc aaa aac ctg gtg act gac			1872
Thr Asn Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn Leu Val Thr Asp			
610	615	620	
cgg gtg aca gag aat atg gcc acg gtc tcc tgg gac ccg gtg cag gcc			1920
Arg Val Thr Glu Asn Met Ala Thr Val Ser Trp Asp Pro Val Gln Ala			
625	630	635	640
gcc att gac aag tac gtg gtg cgc tac acc tct gct ggt gga gag acc			1968
Ala Ile Asp Lys Tyr Val Val Arg Tyr Thr Ser Ala Gly Gly Glu Thr			
645	650	655	
agg gag gtt ccg gtg ggg aag gag cag agc agc aca gtc ctg aca ggc			2016
Arg Glu Val Pro Val Gly Lys Glu Gln Ser Ser Thr Val Leu Thr Gly			
660	665	670	
ctg aga ccg ggt atg gag tac atg gtg cac gtg tgg gcc cag aag ggg			2064
Leu Arg Pro Gly Met Glu Tyr Met Val His Val Trp Ala Gln Lys Gly			
675	680	685	
gac cag gag agc aag aag gcc gac acc aag gcc cag aca gac att gac			
Asp Gln Glu Ser Lys Lys Ala Asp Thr Lys Ala Gln Thr Asp Ile Asp			
690	695	700	
agc ccc caa aac ctg gtg acc gac cgg gtg aca gag aat atg gcc act			2160
Ser Pro Gln Asn Leu Val Thr Asp Arg Val Thr Glu Asn Met Ala Thr			
705	710	715	720
gtc tcc tgg gac ccg gtg cgg gcc acc att gac agg tat gtg gtg cgc			2208
Val Ser Trp Asp Pro Val Arg Ala Thr Ile Asp Arg Tyr Val Val Arg			
725	730	735	
tac acc tct gcc aag gac gga gag acc agg gag gtt ccg gtg ggg aag			2256
Tyr Thr Ser Ala Lys Asp Gly Glu Thr Arg Glu Val Pro Val Gly Lys			
740	745	750	
gag cag agt agc act gtc ctg acg ggc ctg agg ccg ggt gtg gag tac			2304
Glu Gln Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr			
755	760	765	
acg gtg cac gtg tgg gcc cag aag ggg gcc cag gag agc aag aag gct			2352
Thr Val His Val Trp Ala Gln Lys Gly Ala Gln Glu Ser Lys Lys Ala			
770	775	780	

gac acc aag gcc cag aca gac att gac agc ccc caa aac ctg gtc act 2400  
Asp Thr Lys Ala Gln Thr Asp Ile Asp Ser Pro Gln Asn Leu Val Thr  
785 790 795 800

gac tgg gtg aca gag aat aca gcc act gtc tcc tgg gac ccg gtg cag 2448  
Asp Trp Val Thr Glu Asn Thr Ala Thr Val Ser Trp Asp Pro Val Gln  
805 810 815

gcc acc att gac agg tat gtg gtg cac tac acg tct gcc aac gga gag 2496  
Ala Thr Ile Asp Arg Tyr Val Val His Tyr Thr Ser Ala Asn Gly Glu  
820 825 830

acc agg gag gtt cca gtg ggg aag gag cag agc agc act gtc ctg acg 2544  
Thr Arg Glu Val Pro Val Gly Lys Glu Gln Ser Ser Thr Val Leu Thr  
835 840 845

ggc ctg agg ccg ggc atg gag tac acg gtg cac gtg tgg gcc cag aag 2592  
Gly Leu Arg Pro Gly Met Glu Tyr Thr Val His Val Trp Ala Gln Lys  
850 855 860

ggg aac cag gag agc aag aag gct gac acc aag gcc cag gaa att 2640  
Gly Asn Gln Glu Ser Lys Lys Ala Asp Thr Lys Ala Thr Glu Ile  
865 870 875 880

gac ggc ccc aaa aac cta gtg act gac tgg gtg acg gag tgg gcc 2688  
Asp Gly Pro Lys Asn Leu Val Thr Asp Trp Val Thr Glu Asn Met Ala  
885 890 895

act gtc tcc tgg gac ccg gtt cag gcc acc att gac aag tac atg gtg 2736  
Thr Val Ser Trp Asp Pro Val Gln Ala Thr Ile Asp Lys Tyr Met Val  
900 905 910

**cgc tac acc tct gct gac gga gag acc agg gag gtt ccg gtg ggg aag** 2784  
**Arg Tyr Thr Ser Ala Asp Gly Glu Thr Arg Glu Val Pro Val Gly Lys**  
**915 920 925**

gag cac agc agc act gtc ctg acg ggc ctg aga cca ggc atg gag tac 2832  
Glu His Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Met Glu Tyr  
930 935 940

atg gtg cac gtg tgg gcc cag aag ggg gcc cag gag agc aag aag gct 2880  
Met Val His Val Trp Ala Gln Lys Gly Ala Gln Glu Ser Lys Lys Ala  
945 950 955 960

gac acc aag gcc cag aca gaa ctc gac cct ccc aga aac ctt cgt cca 2928  
Asp Thr Lys Ala Gln Thr Glu Leu Asp Pro Pro Arg Asn Leu Arg Pro

965

970

975

tct gct gta acg cag tct ggt ggc ata ttg acc tgg acg ccc ccc tct 2976  
 Ser Ala Val Thr Gln Ser Gly Gly Ile Leu Thr Trp Thr Pro Pro Ser  
 980 985 990

gct cag atc cac ggc tac att ctg act tac cag ttc cca gat ggc aca 3024  
 Ala Gln Ile His Gly Tyr Ile Leu Thr Tyr Gln Phe Pro Asp Gly Thr  
 995 1000 1005

gtt aag gag atg cag ctg gga cgg gaa gac cag agg ttt gcg ttg 3069  
 Val Lys Glu Met Gln Leu Gly Arg Glu Asp Gln Arg Phe Ala Leu  
 1010 1015 1020

caa ggc ctt gag caa ggc gcc acc tac cct gtc tcc ctt gtt gcc 3114  
 Gln Gly Leu Glu Gln Gly Ala Thr Tyr Pro Val Ser Leu Val Ala  
 1025 1030 1035

ttt aag ggt ggt cgc cgg agc aga aat gta tcc acc acc ctc tcc 3159  
 Phe Lys Gly Gly Arg Arg Ser Arg Asn Val Ser Thr Thr Leu Ser  
 1040 1045 1050

aca gtt ggt gcc cgt ttc caa cag cct cgc gac tgc agt cag gtt 3204  
 Thr Val Gly Ala Arg Phe Pro Asn Val Ser Asp Cys Ser Gln Val  
 1055 1060 1065

cag cag aac agc aat gcc gcc agt ggt ctg tac acc atc tac ctg 3249  
 Gln Gln Asn Ser Asn Ala Ala Ser Gly Leu Tyr Thr Ile Tyr Leu  
 1070 1075 1080

cat ggc gat gcc agc cgg ccc ctg cag gtg tac tgt gac atg gaa 3294  
 His Gly Asp Ala Ser Arg Pro Leu Gln Val Tyr Cys Asp Met Glu  
 1085 1090 1095

acg gac gga ggt ggc tgg att gtc ttc cag agg cgg aac act ggg 3339  
 Thr Asp Gly Gly Gly Trp Ile Val Phe Gln Arg Arg Asn Thr Gly  
 1100 1105 1110

cag ctg gat ttc ttc aag cga tgg agg agc tat gtg gaa ggc ttt 3384  
 Gln Leu Asp Phe Phe Lys Arg Trp Arg Ser Tyr Val Glu Gly Phe  
 1115 1120 1125

ggg gac ccc atg aag gag ttc tgg ctt gga ctt gac aag cta cac 3429  
 Gly Asp Pro Met Lys Glu Phe Trp Leu Gly Leu Asp Lys Leu His  
 1130 1135 1140



aac ctc	acc acc ggc act cca	gcg cgg tat gag gtg	aga gtg gat	3474
Asn Leu	Thr Thr Gly Thr Pro	Ala Arg Tyr Glu Val	Arg Val Asp	
1145	1150	1155		
tta cag	act gcc aat gaa tct	gcc tat gct ata tat	gat ttc ttc	3519
Leu Gln	Thr Ala Asn Glu Ser	Ala Tyr Ala Ile Tyr	Asp Phe Phe	
1160	1165	1170		
caa gtg	gcc tcc agc aag gag	cgg tat aag ctg aca	gtt ggg aaa	3564
Gln Val	Ala Ser Ser Lys Glu	Arg Tyr Lys Leu Thr	Val Gly Lys	
1175	1180	1185		
tac aga	ggc acg gca ggg gat	gct ctt act tac cac	aat gga tgg	3609
Tyr Arg	Gly Thr Ala Gly Asp	Ala Leu Thr Tyr His	Asn Gly Trp	
1190	1195	1200		
aag ttt	aca act ttt gac aga	gac aat gat atc gca	ctc agc aac	3654
Lys Phe	Thr Thr Phe Asp Arg	Asp Asn Asp Ile Ala	Leu Ser Asn	
1205	1210	1215		
tgt gcc	aca cat cat ggt	ggc tgg tgg tat aag	aac tgc cac	3699
Cys	Leu Thr His His Gly	Gly Trp Trp Tyr Lys	Asn Cys His	
1220	1225	1230		
ttg gcc	gct cct aat ggc aga	tat ggg gag acc aag	cac agt gag	3744
Leu Ala	Asn Pro Asn Gly Arg	Tyr Gly Glu Thr Lys	His Ser Glu	
1235	1240	1245		
ggg gtg	aac tgg gag cct tgg	aaa gga cat gaa ttc	tcc att cct	3789
Gly Val	Asn Trp Glu Pro Trp	Lys Gly His Glu Phe	Ser Ile Pro	
1250	1255	1260		
tac gtg	gag ttg aaa atc cgc	cct cat ggc tac agc	agg gag cct	3834
Tyr Val	Glu Leu Lys Ile Arg	Pro His Gly Tyr Ser	Arg Glu Pro	
1265	1270	1275		
gtc ctg	ggc aga aag aag cgg	acg ctg aga gga agg	ctg cga acg	3879
Val Leu	Gly Arg Lys Lys Arg	Thr Leu Arg Gly Arg	Leu Arg Thr	
1280	1285	1290		
ttc tga				3885
Phe				

<210> 4  
 <211> 1294  
 <212> PRT  
 <213> homo sapiens

<400> 4

Met Phe Arg Phe Pro Met Gly Leu Leu Leu Gly Ser Val Leu Leu Val  
 1 5 10 15

Ala Ser Ala Pro Ala Thr Leu Glu Pro Pro Gly Cys Ser Asn Lys Glu  
 20 25 30

Gln Gln Val Thr Val Ser His Thr Tyr Lys Ile Asp Val Pro Lys Ser  
 35 40 45

Ala Leu Val Gln Val Asp Ala Asp Pro Gln Pro Leu Ser Asp Asp Gly  
 50 55 60

Ala Ser Leu Leu Ala Leu Gly Glu Ala Arg Glu Glu Gln Asn Ile Ile  
 65 70 75 80

Phe Arg His Asn Ile Arg Leu Gln Thr Pro Gln Lys Asp Cys Glu Leu  
 85 90 95

Ala Gly Ser Val Gln Asp Leu Leu Ala Arg Val Lys Lys Leu Glu Glu  
 100 105 110

Glu Met Val Glu Met Lys Glu Gln Cys Ser Ala Gln Arg Cys Cys Gln  
 115 120 125

Gly Val Thr Asp Leu Ser Arg His Cys Ser Gly His Gly Thr Phe Ser  
 130 135 140

Leu Glu Thr Cys Ser Cys His Cys Glu Glu Gly Arg Glu Gly Pro Ala  
 145 150 155 160

Cys Glu Arg Leu Ala Cys Pro Gly Ala Cys Ser Gly His Gly Arg Cys  
 165 170 175

Val Asp Gly Arg Cys Leu Cys His Glu Pro Tyr Val Gly Ala Asp Cys  
 180 185 190

Gly Tyr Pro Ala Cys Pro Glu Asn Cys Ser Gly His Gly Glu Cys Val  
 195 200 205

Arg Gly Val Cys Gln Cys His Glu Asp Phe Met Ser Glu Asp Cys Ser  
210 215 220

Glu Lys Arg Cys Pro Gly Asp Cys Ser Gly His Gly Phe Cys Asp Thr  
225 230 235 240

Gly Glu Cys Tyr Cys Glu Glu Gly Phe Thr Gly Leu Asp Cys Ala Gln  
245 250 255

Val Val Thr Pro Gln Gly Leu Gln Leu Leu Lys Asn Thr Glu Asp Ser  
260 265 270

Leu Leu Val Ser Trp Glu Pro Ser Ser Gln Val Asp His Tyr Leu Leu  
275 280 285

Ser Tyr Tyr Pro Leu Gly Lys Glu Leu Ser Gly Lys Gln Ile Gln Val  
290 295 300

Pro Lys Glu Gln His Ser Tyr Glu Ile Leu Gly Leu Leu Pro Gly Thr  
305 310 315 320

Lys Tyr Ile Val Thr Leu Arg Asn Val Lys Asn Glu Val Ser Ser Ser  
325 330 335

Pro Gln His Leu Leu Ala Thr Thr Asp Leu Ala Val Leu Gly Thr Ala  
340 345 350

Trp Val Thr Asp Glu Thr Glu Asn Ser Leu Asp Val Glu Trp Glu Asn  
355 360 365

Pro Ser Thr Glu Val Asp Tyr Tyr Lys Leu Arg Tyr Gly Pro Met Thr  
370 375 380

Gly Gln Glu Val Ala Glu Val Thr Val Pro Lys Ser Ser Asp Pro Lys  
385 390 395 400

Ser Arg Tyr Asp Ile Thr Gly Leu His Pro Gly Thr Glu Tyr Lys Ile  
405 410 415

Thr Val Val Pro Met Arg Gly Glu Leu Glu Gly Lys Pro Ile Leu Leu  
420 425 430

Asn Gly Arg Thr Glu Ile Asp Ser Pro Thr Asn Val Val Thr Asp Arg  
435 440 445

Val Thr Glu Asp Thr Ala Thr Val Ser Trp Asp Pro Val Gln Ala Val  
 450 455 460

Ile Asp Lys Tyr Val Val Arg Tyr Thr Ser Ala Asp Gly Asp Thr Lys  
 465 470 475 480

Glu Met Ala Val His Lys Asp Glu Ser Ser Thr Val Leu Thr Gly Leu  
 485 490 495

Lys Pro Gly Glu Ala Tyr Lys Val Tyr Val Trp Ala Glu Arg Gly Asn  
 500 505 510

Gln Gly Ser Lys Lys Ala Asp Thr Asn Ala Leu Thr Glu Ile Asp Ser  
 515 520 525

Pro Ala Asn Leu Val Thr Asp Arg Val Thr Glu Asn Thr Ala Thr Ile  
 530 535 540

Ser Trp Asp Pro Val Gln Ala Thr Ile Asp Lys Tyr Val Val Arg Tyr  
 545 550 555 560

Thr Ser Ala Asp Asp Gln Glu Thr Arg Glu Val Leu Val Gly Lys Glu  
 565 570 575

Gln Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr Thr  
 580 585 590

Val His Val Trp Ala Gln Lys Gly Asp Arg Glu Ser Lys Lys Ala Asp  
 595 600 605

Thr Asn Ala Pro Thr Asp Ile Asp Ser Pro Lys Asn Leu Val Thr Asp  
 610 615 620

Arg Val Thr Glu Asn Met Ala Thr Val Ser Trp Asp Pro Val Gln Ala  
 625 630 635 640

Ala Ile Asp Lys Tyr Val Val Arg Tyr Thr Ser Ala Gly Gly Glu Thr  
 645 650 655

Arg Glu Val Pro Val Gly Lys Glu Gln Ser Ser Thr Val Leu Thr Gly  
 660 665 670

Leu Arg Pro Gly Met Glu Tyr Met Val His Val Trp Ala Gln Lys Gly  
 675 680 685

Asp Gln Glu Ser Lys Lys Ala Asp Thr Lys Ala Gln Thr Asp Ile Asp  
690 695 700

Ser Pro Gln Asn Leu Val Thr Asp Arg Val Thr Glu Asn Met Ala Thr  
705 710 715 720

Val Ser Trp Asp Pro Val Arg Ala Thr Ile Asp Arg Tyr Val Val Arg  
725 730 735

Tyr Thr Ser Ala Lys Asp Gly Glu Thr Arg Glu Val Pro Val Gly Lys  
740 745 750

Glu Gln Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Val Glu Tyr  
755 760 765

Thr Val His Val Trp Ala Gln Lys Gly Ala Gln Glu Ser Lys Lys Ala  
770 775 780

Asp Thr Lys Ala Gln Thr Asp Ile Asp Ser Pro Gln Asn Leu Val Thr  
785 790 795 800

Asp Trp Val Thr Glu Asn Thr Ala Thr Val Ser Trp Asp Pro Val Gln  
805 810 815

Ala Thr Ile Asp Arg Tyr Val Val His Tyr Thr Ser Ala Asn Gly Glu  
820 825 830

Thr Arg Glu Val Pro Val Gly Lys Glu Gln Ser Ser Thr Val Leu Thr  
835 840 845

Gly Leu Arg Pro Gly Met Glu Tyr Thr Val His Val Trp Ala Gln Lys  
850 855 860

Gly Asn Gln Glu Ser Lys Lys Ala Asp Thr Lys Ala Gln Thr Glu Ile  
865 870 875 880

Asp Gly Pro Lys Asn Leu Val Thr Asp Trp Val Thr Glu Asn Met Ala  
885 890 895

Thr Val Ser Trp Asp Pro Val Gln Ala Thr Ile Asp Lys Tyr Met Val  
900 905 910

Arg Tyr Thr Ser Ala Asp Gly Glu Thr Arg Glu Val Pro Val Gly Lys  
915 920 925

Glu His Ser Ser Thr Val Leu Thr Gly Leu Arg Pro Gly Met Glu Tyr  
 930 935 940

Met Val His Val Trp Ala Gln Lys Gly Ala Gln Glu Ser Lys Lys Ala  
 945 950 955 960

Asp Thr Lys Ala Gln Thr Glu Leu Asp Pro Pro Arg Asn Leu Arg Pro  
 965 970 975

Ser Ala Val Thr Gln Ser Gly Gly Ile Leu Thr Trp Thr Pro Pro Ser  
 980 985 990

Ala Gln Ile His Gly Tyr Ile Leu Thr Tyr Gln Phe Pro Asp Gly Thr  
 995 1000 1005

Val Lys Glu Met Gln Leu Gly Arg Glu Asp Gln Arg Phe Ala Leu  
 1010 1015 1020

Gln Gly Leu Glu Gln Gly Ala Thr Tyr Pro Val Ser Leu Val Ala  
 1025 1030 1035

Phe Lys Gly Gly Arg Arg Ser Arg Asn Val Ser Thr Thr Leu Ser  
 1040 1045 1050

Thr Val Gly Ala Arg Phe Pro His Pro Ser Asp Cys Ser Gln Val  
 1055 1060 1065

Gln Gln Asn Ser Asn Ala Ala Ser Gly Leu Tyr Thr Ile Tyr Leu  
 1070 1075 1080

His Gly Asp Ala Ser Arg Pro Leu Gln Val Tyr Cys Asp Met Glu  
 1085 1090 1095

Thr Asp Gly Gly Gly Trp Ile Val Phe Gln Arg Arg Asn Thr Gly  
 1100 1105 1110

Gln Leu Asp Phe Phe Lys Arg Trp Arg Ser Tyr Val Glu Gly Phe  
 1115 1120 1125

Gly Asp Pro Met Lys Glu Phe Trp Leu Gly Leu Asp Lys Leu His  
 1130 1135 1140

Asn Leu Thr Thr Gly Thr Pro Ala Arg Tyr Glu Val Arg Val Asp  
 1145 1150 1155

Leu Gln Thr Ala Asn Glu Ser Ala Tyr Ala Ile Tyr Asp Phe Phe  
 1160 1165 1170

Gln Val Ala Ser Ser Lys Glu Arg Tyr Lys Leu Thr Val Gly Lys  
 1175 1180 1185

Tyr Arg Gly Thr Ala Gly Asp Ala Leu Thr Tyr His Asn Gly Trp  
 1190 1195 1200

Lys Phe Thr Thr Phe Asp Arg Asp Asn Asp Ile Ala Leu Ser Asn  
 1205 1210 1215

Cys Ala Leu Thr His His Gly Gly Trp Trp Tyr Lys Asn Cys His  
 1220 1225 1230

Leu Ala Asn Pro Asn Gly Arg Tyr Gly Glu Thr Lys His Ser Glu  
 1235 1240 1245

Gly Val Asn Trp Glu Pro Trp Lys Gly His Glu Phe Ser Ile Pro  
 1250 1255 1260

Tyr Val Glu Leu Lys Ile Arg Pro His Gly Tyr Ser Arg Glu Pro  
 1265 1270 1275

Val Leu Gly Arg Lys Lys Arg Thr Leu Arg Gly Arg Leu Arg Thr  
 1280 1285 1290

Phe

<210> 5  
 <211> 25  
 <212> DNA  
 <213> homo sapiens

<400> 5  
 tagcagccca cagcatctac ttgcc

25

<210> 6  
 <211> 27  
 <212> DNA  
 <213> homo sapiens

<400> 6  
attgctgttc tgctgaacct gactgca 27

<210> 7  
<211> 34  
<212> DNA  
<213> synthetic

<400> 7  
atggatccag aaattgacgg ccccaaaaac ctag 34

<210> 8  
<211> 33  
<212> DNA  
<213> synthetic

<400> 8  
ataagcttgt ggagaggggtg gtggatacat cc 33

<210> 9  
<211> 24  
<212> DNA  
<213> Mus musculus

<400> 9  
aggagatggt ggctgtattt tcgg 24

<210> 10  
<211> 24  
<212> DNA  
<213> Saccharomyces cerevisiae vector pJG4-5

<400> 10  
agcctcttgc tgagtggaga tgcc 24

<210> 11  
<211> 30  
<212> DNA



<213> synthetic

<400> 11

tagaattcgg tcacctgatt ggtcactagg

30

<210> 12

<211> 22

<212> DNA

<213> Saccharomyces cerevisiae vector pJG4-5

<400> 12

ttatgatgtg ccagattatg cc

22

<210> 13

<211> 26

<212> DNA

<213> Mus musculus

<400> 13

ctcaaattga cctcaccacc cctacc

26

<210> 14

<211> 24

<212> DNA

<213> Saccharomyces cerevisiae vector pJG4-5

<400> 14

aagccgacaa ccttgattgg agac

24

<210> 15

<211> 22

<212> DNA

<213> Mus musculus

<400> 15

taccagttcc caaatggcac cg

22

<210> 16

<211> 21

<212> DNA

<213> Saccharomyces cerevisiae vector pJG4-5

<400> 16  
aaacctctgg cgaagaagtc c

21

<210> 17  
<211> 25  
<212> DNA  
<213> synthetic

<400> 17  
catcctggag ggtctgctcc

20

<210> 18  
<211> 20  
<212> DNA  
<213> synthetic

<400> 18  
gggcattggg gtcagctttc

20

<210> 19  
<211> 28  
<212> DNA  
<213> synthetic

<400> 19  
gactcgagct ttccaaggat gagtctcc

28

<210> 20  
<211> 26  
<212> DNA  
<213> synthetic

<400> 20  
gaggatcccc tggttgcccc ttccag

26

<210> 21  
<211> 20  
<212> DNA  
<213> synthetic

<400> 21  
gcgctacact tctgctgatg 20

<210> 22  
<211> 18  
<212> DNA  
<213> synthetic  
<400> 22  
ctgtggagag ggtggtgg 18

<210> 23  
<211> 27  
<212> DNA  
<213> synthetic  
<400> 23  
gactcgagtg cacaaggatg agagcag 27

<210> 24  
<211> 28  
<212> DNA  
<213> synthetic  
<400> 24  
gaggatccac ccttaaaggc aacaaggg 28

<210> 25  
<211> 28  
<212> DNA  
<213> synthetic  
<400> 25  
cgcagtctgg tggcatattg 20

<210> 26  
<211> 19  
<212> DNA  
<213> synthetic  
<400> 26  
catgatttgt tctgcgggc 19

<210> 27  
<211> 28  
<212> DNA  
<213> synthtetic

<400> 27  
gactcgagcg gctacattct gacttacc

28

<210> 28  
<211> 29  
<212> DNA  
<213> synthetic

<400> 28  
gaggatcctc agtgatggtg atggtgatg

29

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**